

ON POWERS OF COVER IDEALS OF GRAPHS

DANCHENG LU AND ZEXIN WANG

ABSTRACT. For a simple graph G , assume that $J(G)$ is the vertex cover ideal of G and $J(G)^{(s)}$ is the s -th symbolic power of $J(G)$. We prove that $\text{reg}(J(G)^{(s)}) = \text{reg}(J(G)^s)$ for all $s \geq 1$ and for all odd cycle C . For a simplicial complex Δ , we show that if I_Δ^\vee is weakly polymatroidal (not necessarily generated in one degree) then Δ is vertex decomposable. Some evidences are provided that the converse conclusion of the above result also holds true if Δ is pure. Let $W = G^\pi$ be a fully clique-whiskering graph. We prove that $J(W)^s$ is weakly polymatroidal for all $s \geq 1$.

1. INTRODUCTION

Let $R = K[x_1, \dots, x_n]$ be the polynomial ring over a field K and let G be a simple graph on vertex set $[n] := \{1, 2, \dots, n\}$ with edge set $E(G)$. There are two square-free monomial ideals of R associated to G : the *edge ideal* $I(G)$ which is generated by all monomial $x_i x_j$ with $\{i, j\} \in E(G)$ and the *vertex cover ideal* $J(G)$ generated by monomials $\prod_{i \in F} x_i$, where F is taken over all minimal vertex covers of G . Recall that a subset F of $V(G)$ is a *vertex cover* of G if $F \cap e \neq \emptyset$ for every edge e of G and a vertex cover F of G is *minimal* if $F \setminus \{i\}$ is not a vertex cover for each $i \in F$. The vertex cover ideal $J(G)$ is the Alexander dual of the edge ideal $I(G)$, i.e.,

$$J(G) = I(G)^\vee = \bigcap_{\{i,j\} \in E(G)} (x_i, x_j).$$

Let I be a graded ideal of R . The s -th *symbolic power* of I is defined by

$$I^{(s)} = \bigcap_{\mathfrak{p} \in \text{Min}(I)} I^s R_{\mathfrak{p}} \cap R,$$

where $\text{Min}(I)$ is as usual the set of all minimal prime ideals of I . It follows from [12, Proposition 1.4.4] that for every integer $s \geq 1$,

$$J(G)^{(s)} = \bigcap_{\{i,j\} \in E(G)} (x_i, x_j)^s.$$

The Castelnuovo-Mumford regularity (or simply regularity) is a fundamental invariant in commutative algebra and algebraic geometry. For a finitely generated graded module M over the polynomial ring R , the *regularity* of M , denoted by $\text{reg}(M)$, is the least integer $r \geq 0$ such that for all $i \geq 0$, the i -th syzygy of M is

2010 *Mathematics Subject Classification.* Primary 13C70, 13D45 Secondary 05E40.

Key words and phrases. vertex cover ideal, regularity, symbolic power, vertex decomposable, odd cycle, weakly polymatroidal, whisker graph.

generated by homogeneous elements of degree at most $r+i$. An equivalent definition of the regularity via local cohomology is as follows:

$$\text{reg}(M) = \max\{i + j : H_{\mathfrak{m}}^i(M)_j \neq 0, i \geq 0, j \in \mathbb{Z}\}.$$

Here \mathfrak{m} denotes the maximal ideal (x_1, \dots, x_n) .

For edge ideals of graphs, there have been a lot of research on connections between the regularity functions $\text{reg}(I(G)^s)$ as well as $\text{reg}(I(G)^{(s)})$ and the combinatorial properties of G , see [1] and the references therein. Recently the conjecture that $\text{reg}(I(G)^{(s)}) = \text{reg}(I(G)^s)$ for all $s \geq 1$ and for all graphs G attracted much attention and much progress has been made in this direction. For the details, see [9] and the references therein.

Meanwhile, the study of algebraic properties of (symbolic and ordinary) powers of vertex cover ideals of graphs is also an active research topic. However, the regularity of powers of such ideals is harder to compute or deal with. In fact, although S.A. Seyed Fakhari presented in [23] the following remarkable bounds for a large class of graphs G , including bipartite graphs, unmixed graphs, claw-free graphs:

$$s\text{Deg}(J(G)) \leq \text{reg}(J(G)^{(s)}) \leq (s-1)\text{Deg}(J(G)) + |V(G)| - 1,$$

there are not many graphs G for which either $\text{reg}(J(G)^{(s)})$ or $\text{reg}(J(G)^s)$ is known precisely. Here, $\text{Deg}(J(G))$ is the maximum size of minimal vertex covers of G . When G is either a crown graph or a complete multipartite graph, $\text{reg}(J(G)^{(s)})$ was explicitly given in [11]. On the other side, if a graded ideal is componentwise linear then its regularity is equal to the maximum degree of its minimal generators. In the literatures [4, 7, 16, 23, 21] and [24], some classes of graphs for which either $J(G)^s$ or $J(G)^{(s)}$ is componentwise linear are identified. For examples, it was proved in [24] that G is a Cohen-Macaulay very well-covered graph if and only if $J(G)^{(s)}$ has a linear resolution for some (equivalently, for all) integer $s \geq 2$. In [4, 7, 21, 23] and [24], among others, they investigate the question of how to combinatorially modify a graph to obtain componentwise linearity of the corresponding monomial ideals, and identify many graphs G such that $J(G)^{(s)}$ is componentwise linear. In [16], it was proved that if G is a Cohen-Macaulay cactus graph then $J(G)^s$ has a linear resolution for all $s \geq 1$. For such graphs G , the regularity of either $J(G)^s$ or $J(G)^{(s)}$ is known. More recently, a new upper bound for $\text{reg}(J(G)^{(s)})$ was given in [19] when G is a non-bipartite graph.

In this paper we investigate further the properties of (symbolic and ordinary) powers of vertex cover ideals of simple graphs. Our first main result is motivated by Theorem 5.15 in [4], in which a family of graphs G was constructed such that $\text{reg}(J(G)^{(s)})$ is not eventually linear in s . This result particularly shows that the equality $\text{reg}(J(G)^{(s)}) = \text{reg}(J(G)^s)$ is not true in general. On the other side, we have known the equality $\text{reg}(I(G)^{(s)}) = \text{reg}(I(G)^s)$ for all $s \geq 1$ holds for many classes of graphs such as unicyclic graphs and chordal graphs and so on. These facts lead us to ask the following question:

For which graphs G has one $\text{reg}(J(G)^{(s)}) = \text{reg}(J(G)^s)$ for all $s \geq 1$?

In this vein we prove the following result.

Theorem 2.4 *If C is an odd cycle, then $\text{reg}(J(C)^{(s)}) = \text{reg}(J(C)^s)$ for all $s \geq 1$.*

We remark this is the first non-trivial example where the above formula holds, due to the well-known fact that G is a bipartite graph if and only if $J(G)^{(s)} = J(G)^s$ for some $s \geq 2$ (equivalently for all $s \geq 1$), see [25, Proposition 1.3] or [13, Theorem 5.1].

We next investigate relations between weak polymatroidality of a monomial ideal and vertex decomposability of a simplicial complex. It turns out it helps to understand the behaviors of the powers of cover ideals. Vertex decomposability was first introduced by [20] in the pure case, and extended to the non-pure case in [2]. It is defined in terms of the deletion and link. Let Δ be a simplicial complex on vertex set V . For $x \in V$, the *link* of x in Δ is the subcomplex

$$\text{lk}_\Delta(x) = \{F \in \Delta: F \cup \{x\} \in \Delta \text{ and } x \notin F\}$$

and the *deletion* of x in Δ is the subcomplex

$$\text{del}_\Delta(x) = \{F \in \Delta: x \notin F\}.$$

Definition 1.1. A simplicial complex Δ is said to be *vertex decomposable* if either Δ is a simplex, or there exists a vertex x of Δ such that

- (1) $\text{lk}_\Delta(x)$ and $\text{del}_\Delta(x)$ are vertex decomposable;
- (2) Each facet of $\text{lk}_\Delta(x)$ is not a facet of $\text{del}_\Delta(x)$.

A vertex satisfying condition (2) is called a *shedding vertex* of Δ .

For the recent developments on vertex decomposability, one may refer to [8] and the references therein. The following strict implications is well-known for a simplicial complex Δ :

$$\text{vertex decomposable} \implies \text{shellable} \implies \text{sequentially Cohen-Macaulay}$$

Moreover, Δ is shellable if and only if I_Δ^\vee has linear quotients; and Δ is sequentially Cohen-Macaulay if and only if I_Δ^\vee is componentwise linear. One may ask what property I_Δ^\vee has when Δ is vertex decomposable, or vice versa.

Definition 1.2. Following [10], we say that a monomial ideal I in R is *weakly polymatroidal* if for every pair of elements $u = x_1^{a_1} \cdots x_n^{a_n}$ and $v = x_1^{b_1} \cdots x_n^{b_n}$ of $G(I)$ with $a_1 = b_1, \dots, a_{q-1} = b_{q-1}$ and $a_q < b_q$, (noting that $q < n$) there exists $p > q$ such that $w := (x_q u)/x_p$ belongs to $G(I)$. Here, $G(I)$ denotes the set of minimal monomial generators of I .

Different from the original definition in [10], we here do not require I to be generated in one degree. Using the same method as in [10], one can prove if I is weakly polymatroidal in this generalized sense, then I has linear quotients. Our second main result is as follows:

Theorem 3.1. *If I_Δ^\vee is weakly polymatroidal then Δ is vertex decomposable.*

This particularly shows that being weakly polymatroidal is a condition stronger than the property of having linear quotients for a monomial ideal. We also prove that the converse of Theorem 3.1 holds in some special cases. Recall that a graph G is *unmixed* if every minimal vertex cover of G has the same cardinality, i.e. $J(G)$

is generated in one degree, and that G is vertex decomposable if the independence complex of G is vertex decomposable.

Corollary 3.2. *Let G be either a cactus graph or a bipartite graphs or a chordal graph. Assume further that G is unmixed. Then the following are equivalent:*

- (1) $R/I(G)$ is Cohen-Macaulay;
- (2) G is vertex decomposable;
- (3) $J(G)$ is weakly polymatroidal (in some ordering of variables);
- (4) $J(G)^s$ is weakly polymatroidal (in some ordering of variables) for all $s \geq 1$.

Here, a *cactus* graph is a simple graph in which every edge belongs to at most one cycle. This leads us to conjecture the following.

Conjecture 2. *If G is an unmixed graph, then G is vertex decomposable if and only if $J(G)$ is weakly polymatroidal.*

The condition that G is unmixed cannot be dropped in the above conjecture. We show that the above conjecture holds true if either G has girth ≥ 5 or G is very well-covered, see Propositions 3.6 and 3.7.

It is natural to ask for which unmixed graphs G , $J(G)^s$ are weakly polymatroidal besides the graphs given in Corollary 3.2. Of course, such graphs (i.e., $R/I(G)$) should be Cohen-Macaulay by Alexander duality. Let G be a simple graph on vertex set $V(G)$ with edge set $E(G)$. Following [3], a *clique vertex-partition* of G is a partition $V(G) = W_1 \sqcup \cdots \sqcup W_t$ such that the induced graph of G on W_i is a clique (a complete graph) for $i = 1, \dots, t$. Denote this partition by $\pi = \{W_1, \dots, W_t\}$. The *fully clique-whiskering graph* G^π of G by π is the graph on vertex set $V(G) \cup \{y_1, \dots, y_t\}$ and with edge set $E(G) \cup \{vy_i : v \in W_i, 1 \leq i \leq t\}$. When π is a trivial partition, i.e., $|W_1| = \cdots = |W_t| = 1$, G^π is also called the *whisker graph* of G . A tree is Cohen-Macaulay if and only if it is the whisker graph of some other tree. A Cameron-Walker graph is Cohen-Macaulay if and only if it is a fully clique-whiskering graph G^π of a bipartite graph G by some clique vertex-partition π of G , see [14, Theorem 1.3]. Our third main result is as follows:

Theorem 4.3. *If $W = G^\pi$ for some graph G and some clique vertex-partition π , then $J(W)^s$ is weakly polymatroidal for all $s \geq 1$.*

This result is a complement of Corollary 3.2. As a consequence, we obtain if $W = G^\pi$ then $\text{reg}(J(W)^s) = \text{reg}(J(W)^{(s)}) = s|V(G)|$ for all $s \geq 1$. The following is another consequence of Theorem 4.3.

Corollary 4.5. *If W is the whisker graph of some graph, then both $J(W)^s$ and $J(W)^{(s)}$ are weakly polymatroidal for all $s \geq 1$.*

In the rest part of this paper we will keep the notions introduced in this section unless otherwise stated, and refer to [12] for some unexplained notions.

2. POWERS OF COVER IDEALS OF ODD CYCLES

In this section we will prove that if C is an odd cycle then both $J(C)^s$ and $J(C)^{(s)}$ have the same regularity for all $s \geq 1$.

We begin with fixing some notions. Let M be a finitely generated graded R -module generated by homogeneous elements f_1, \dots, f_r minimally with $\deg(f_1) \leq$

$\deg(f_2) \leq \dots \leq \deg(f_r)$. We denote by $\text{Deg}(M)$ the number $\deg(f_r)$ and by $\deg(M)$ the number $\deg(f_1)$. It is known that $\text{reg}(M) \geq \text{Deg}(M)$. Let \mathfrak{m} denote the maximal graded ideal (x_1, x_2, \dots, x_n) of R .

Proposition 2.1. *Let I be a homogeneous ideal in R and t a positive integer. Put $J = I \cap \mathfrak{m}^t$. Then the following holds.*

- (1) $\text{reg}(\frac{R}{I}) \leq \text{reg}(\frac{R}{J})$;
- (2) If $t \leq \text{Deg}(I)$, then $\text{reg}(\frac{R}{I}) = \text{reg}(\frac{R}{J})$.

Proof. (1) Set $b_i = \max\{j: H_{\mathfrak{m}}^i(\frac{R}{I})_j \neq 0\}$ and $a_i = \max\{j: H_{\mathfrak{m}}^i(\frac{R}{J})_j \neq 0\}$. Then $\text{reg}(\frac{R}{I}) = \max\{b_i + i: i \geq 0\}$ and $\text{reg}(\frac{R}{J}) = \max\{a_i + i: i \geq 0\}$. Applying the local cohomological functors with respect to \mathfrak{m} to the short exact sequence

$$(\dagger) \quad 0 \rightarrow \frac{R}{J} \rightarrow \frac{R}{I} \oplus \frac{R}{\mathfrak{m}^t} \rightarrow \frac{R}{\mathfrak{m}^t + I} \rightarrow 0,$$

we obtain the long exact sequence: $0 \rightarrow H_{\mathfrak{m}}^0(R/J) \rightarrow H_{\mathfrak{m}}^0(\frac{R}{I}) \oplus H_{\mathfrak{m}}^0(\frac{R}{\mathfrak{m}^t}) \rightarrow H_{\mathfrak{m}}^0(\frac{R}{\mathfrak{m}^t + I}) \rightarrow \dots \rightarrow H_{\mathfrak{m}}^i(R/J) \rightarrow H_{\mathfrak{m}}^i(\frac{R}{I}) \oplus H_{\mathfrak{m}}^i(\frac{R}{\mathfrak{m}^t}) \rightarrow H_{\mathfrak{m}}^i(\frac{R}{\mathfrak{m}^t + I}) \rightarrow \dots$. From this sequence as well as the equality $H_{\mathfrak{m}}^i(\frac{R}{\mathfrak{m}^t + I}) = H_{\mathfrak{m}}^i(\frac{R}{\mathfrak{m}^t}) = 0$ for all $i > 0$, we obtain the following facts.

- (i) $a_i = b_i$ for all $i \geq 2$.
- (ii) The sequence $H_{\mathfrak{m}}^1(\frac{R}{J}) \rightarrow H_{\mathfrak{m}}^1(\frac{R}{I}) \rightarrow 0$ is exact. From this, we have $H_{\mathfrak{m}}^1(\frac{R}{I})_i = 0$ if $H_{\mathfrak{m}}^1(\frac{R}{J})_i = 0$. This implies that $a_1 \geq b_1$.
- (iii) The sequence $0 \rightarrow H_{\mathfrak{m}}^0(\frac{R}{J}) \rightarrow H_{\mathfrak{m}}^0(\frac{R}{I}) \oplus \frac{R}{\mathfrak{m}^t} \rightarrow \frac{R}{I + \mathfrak{m}^t}$ is exact. Thus, for any $i \in \mathbb{Z}$ with $H_{\mathfrak{m}}^0(\frac{R}{J})_i = 0$, we have $\dim_k H_{\mathfrak{m}}^0(\frac{R}{I})_i + \dim_k [\frac{R}{\mathfrak{m}^t}]_i \leq \dim_k [\frac{R}{I + \mathfrak{m}^t}]_i \leq \dim_k [\frac{R}{\mathfrak{m}^t}]_i$ and so $\dim_k H_{\mathfrak{m}}^0(\frac{R}{I})_i = 0$. Hence $a_0 \geq b_0$.

Combining (i),(ii) with (iii), we obtain that $\text{reg}(\frac{R}{I}) \geq \text{reg}(\frac{R}{J})$.

(2) By using the short exact sequence (\dagger) , we also obtain

$$\text{reg}(\frac{R}{J}) \leq \max\{\text{reg}(\frac{R}{I}), t - 1\} = \text{reg}(\frac{R}{I}).$$

This finishes the proof. \square

Proposition 2.1 can be extended to the case of graded modules. Let $M = \bigoplus_{i \in \mathbb{Z}} M_i$ be a finitely generated \mathbb{Z} -graded R -module. For $j \in \mathbb{Z}$, we denote by $M_{\geq j}$ the graded submodule $\bigoplus_{i \geq j} M_i$ of M . Note that $I \cap \mathfrak{m}^t = I_{\geq t}$ for any graded ideal I , we may look upon the following result as a generalization of Proposition 2.1.

Proposition 2.2. *Let M be a finitely generated \mathbb{Z} -graded R -module and let $j \in \mathbb{Z}$ such that $M_{\geq j} \neq 0$. Then the following statements hold:*

- (1) $\text{reg}(M_{\geq j}) \geq \text{reg}(M)$;
- (2) If $j \leq \text{Deg}(M)$, then $\text{reg}(M_{\geq j}) = \text{reg}(M)$.

Proof. (1) Consider the following short exact sequence:

$$(\ddagger) \quad 0 \rightarrow M_{\geq j} \rightarrow M \rightarrow M/M_{\geq j} \rightarrow 0.$$

It is known that $\text{reg}(N) = \max\{i: N_i \neq 0\}$ whenever N is a graded R -module of finite length. From this, it follows that $\text{reg}(M_{\geq j}) \geq j > \text{reg}(M/M_{\geq j})$. Hence $\text{reg}(M) \leq \max\{\text{reg}(M_{\geq j}), \text{reg}(M/M_{\geq j})\} = \text{reg}(M_{\geq j})$.

(2) Using the sequence (‡) again, we obtain

$$\text{reg}(M_{\geq j}) \leq \max\{\text{reg}(M), \text{reg}(M/M_{\geq j}) + 1\}.$$

Note that $\text{reg}(M) \geq \text{Deg}(M) \geq j \geq \text{reg}(M/M_{\geq j}) + 1$, the result follows. \square

Let G be a simple graph on vertex set $[n]$ and H a subgraph of G . The neighborhood of H is defined by

$$N_G(H) = \{i \in V(G): i \text{ is adjacent to some vertex of } H\}.$$

By [13, Proposition 5.3], if $N_G(C) = [n]$ for every odd cycle C , then the symbolic Rees algebra

$$\mathcal{R}_s(J(G)) = \bigoplus_{k \geq 0} J(G)^{(k)} t^k$$

of $J(G)$ is generated by the monomial $x_1 \cdots x_n t^2$ together with the monomials $t \prod_{i \in F} x_i$ such that F is a minimal vertex cover of G . Thus, the following result is a direct consequence of [13, Proposition 5.3].

Proposition 2.3. *Let G be a simple graph on vertex set $[n]$ such that $N_G(C) = [n]$ for every odd cycle of G . Then*

$$J(G)^{(s)} = J(G)^s + \sum_{i=1}^{\lfloor \frac{s}{2} \rfloor} (x_1 x_2 \cdots x_n)^i J(G)^{s-2i}.$$

Here, $\lfloor \frac{s}{2} \rfloor$ denotes the largest integer at most $\frac{s}{2}$.

Let C be an odd cycle of length $n = 2r + 1$. It is not difficult to see that C is not unmixed if $n \geq 9$. More precisely, we have $\text{deg}(J(C)) = r + 1$ and

$$\text{Deg}(J(C)) = \begin{cases} 4t + 2, & n = 6t + 3; \\ 4t + 3, & n = 6t + 5; \\ 4t + 4, & n = 6t + 7. \end{cases}$$

for all $t \geq 0$.

We now come to the main result of this section.

Theorem 2.4. *Let J be the vertex cover ideal of an odd cycle of C on vertex set $[n]$ with $n = 2r + 1$. Then $\text{reg}(J^{(s)}) = \text{reg}(J^s)$ for all $s \geq 1$.*

Proof. Put $t = \text{Deg}(J)$. Then $t \geq r + 1$ and so $\text{Deg}(J^{(s)}) = \text{Deg}(J^s) = st$ for all $s \geq 1$ by Proposition 2.3. Fix $s \geq 1$. We claim that

$$J^{(s)} \cap \mathfrak{m}^{st} = J^s \cap \mathfrak{m}^{st}.$$

In light of Proposition 2.3, it suffices to show that

$$(x_1 x_2 \cdots x_{2r+1})^i J^{s-2i} \cap \mathfrak{m}^{st} \subseteq J^s$$

for all $i = 1, \dots, \lfloor \frac{s}{2} \rfloor$. Fix $i \geq 1$ and let α be a monomial in $(x_1 x_2 \cdots x_{2r+1})^i J^{s-2i} \cap \mathfrak{m}^{st}$. Then we may write $\alpha = (x_1 x_2 \cdots x_{2r+1})^i \alpha_1 \cdots \alpha_{s-2i} \mathbf{u}$, where α_i is a minimal monomial generator of J for each i and \mathbf{u} is some monomial. Since $\text{deg}(\alpha) \geq st$ and $\text{deg}(\alpha_i) \leq$

t , it follows that $\deg(\mathbf{u}) \geq i$. This, together with the fact $(x_1 x_2 \cdots x_{2r+1})x_i \in J^2$ for any $i \in [2r+1]$, implies $\alpha \in J^s$. Thus, the claim is proved.

In view of Proposition 2.1, the result is immediate from the above claim. \square

Remark 2.5. Proposition 2.1 is also useful in the study of powers of edge ideals. For example, [9, Theorem 3.5] is a direct consequence of [9, Corollary 3.3] in view of Proposition 2.1.

3. VERTEX DECOMPOSABILITY VIA WEAK POLYMATROIDALITY

Let Δ be a simplicial complex on $[n]$. In this section we prove that if I_Δ^\vee is weakly polymatroidal then Δ is vertex decomposable. This particularly shows that for a monomial ideal, the property of being weakly polymatroidal is stronger than the property of having linear quotients. The converse implication of the above result is also discussed.

We first give an observation on the property of a shedding vertex. Let Δ be a simplicial complex on vertex set $[n]$ with facets F_1, F_2, \dots, F_r . Assume that k is a shedding vertex of Δ and assume without loss of generality that $k \in F_i$ for $i = 1, \dots, j$ and $k \notin F_i$ for $i = j+1, \dots, r$. Then $\text{lk}_\Delta(k) = \langle F_1 \setminus \{k\}, \dots, F_j \setminus \{k\} \rangle$ and $\text{del}_\Delta(k) = \langle F_1 \setminus \{k\}, \dots, F_j \setminus \{k\}, F_{j+1}, \dots, F_r \rangle = \langle F_{j+1}, \dots, F_r \rangle$. This observation is useful in the following proofs. We also need some more notation. Let I be a monomial ideal. As usual, $G(I)$ denotes the set of minimal monomial generators of I and $\text{supp}(I)$ is the set $\cup_{u \in G(I)} \text{supp}(u)$, where for a monomial u , $\text{supp}(u)$ denotes the set $\{i \in [n] : x_i | u\}$. For a subset $A \subseteq [n]$, $x_{\bar{A}}$ denotes the monomial $\prod_{i \in [n] \setminus A} x_i$.

Theorem 3.1. *Let Δ be a simplicial complex on $[n]$ and suppose that I_Δ^\vee is weakly polymatroidal in some ordering of variables. Then Δ is vertex decomposable.*

Proof. Since the vertex decomposability of a simplicial complex is independent of the ordering of variables, we may assume I_Δ^\vee is weakly polymatroidal itself. Let $\mathcal{F}(\Delta)$ denote the set of facets of Δ . In the following, we will use the induction on $|\mathcal{F}(\Delta)|$, the number of facets of Δ . If $|\mathcal{F}(\Delta)| = 1$, then I_Δ^\vee is generated by a single monomial and so it is weakly polymatroidal automatically. Now assume $|\mathcal{F}(\Delta)| \geq 2$. In this case we let $k = \min\{i : i \in F_1 \cup \dots \cup F_r, i \notin F_1 \cap \dots \cap F_r\}$.

We first show that k is a shedding vertex of Δ . For this, let F, G be facets of Δ such that $k \in F$ and $k \notin G$, respectively. Note that we may write $x_{\bar{F}} = x_{k+1}^{a_{k+1}} \cdots x_n^{a_n}$ and $x_{\bar{G}} = x_k x_{k+1}^{b_{k+1}} \cdots x_n^{b_n}$, where $a_i, b_i \in \{0, 1\}$ for all i and they are minimal generators of I_Δ^\vee . This implies there exists $\ell > k$ such that $\mathbf{u} := x_k x_{\bar{F}} / x_\ell$ is also a minimal generator of I_Δ^\vee and so there exists a facet H of Δ such that $\mathbf{u} = x_{\bar{H}}$. From this it follows that $k \notin H$ and $F \setminus \{k\} \subseteq H$. Since H is a facet of Δ , we have $F \setminus \{k\} \subsetneq H$. This actually shows that none of facets of $\text{lk}_\Delta(k)$ is a facet of $\text{del}_\Delta(k)$ and it follows that k is a shedding vertex of Δ .

Set $\Delta_1 := \text{lk}_\Delta(k)$ and $\Delta_2 := \text{del}_\Delta(k)$. Next we show $I_{\Delta_1}^\vee$ and $I_{\Delta_2}^\vee$ are both weakly polymatroidal. We will look Δ_1 and Δ_2 upon as simplicial complexes on $V := [n] \setminus \{k\}$. Then $I_{\Delta_i}^\vee = (x_{V \setminus F} : F \in \mathcal{F}(\Delta_i))$. Note that

$$\clubsuit \quad G(I_\Delta^\vee) = G(I_{\Delta_1}^\vee) \bigsqcup \{x_k u : u \in G(I_{\Delta_2}^\vee)\}.$$

Moreover, $\text{supp}(I_{\Delta_i}^\vee) \subseteq \{k+1, \dots, n\}$ for $i = 1, 2$.

Let $\mathbf{u} = x_{k+1}^{a_{k+1}} \cdots x_n^{a_n}$ and $\mathbf{v} = x_{k+1}^{b_{k+1}} \cdots x_n^{b_n}$ be distinct elements in $G(I_{\Delta_1}^\vee)$ such that $a_{k+1} = b_{k+1}, \dots, a_{k+i-1} = b_{k+i-1}$ and $a_{k+i} < b_{k+i}$. Then, since I_{Δ}^\vee is weakly polymatroidal, there exists $j > i \geq 1$ such that $\mathbf{w} := x_{k+i}\mathbf{u}/x_{k+j} \in G(I_{\Delta}^\vee)$. This, together with the decomposition in (\clubsuit) , implies $\mathbf{w} \in G(I_{\Delta_1}^\vee)$. Thus, we have proven that $I_{\Delta_1}^\vee$ is weakly polymatroidal. Similarly, we can prove $I_{\Delta_2}^\vee$ is weakly polymatroidal. By induction hypothesis, we have Δ_i is vertex decomposable for $i = 1, 2$ and so Δ is vertex decomposable, as required. \square

Let G be a simple graph. A subset A of $V(G)$ is an *independent set* of G if for any $i, j \in A$, the pair $\{i, j\}$ is not an edge of G . The *independence complex* of G is the collection of all independent sets of G . We call G to be vertex decomposable if its independence complex is vertex decomposable. If we let Δ be the independence complex of G , then the Stanley-Reisner ideal I_Δ is the edge ideal $I(G)$ and its Alexander dual I_Δ^\vee is the vertex cover ideal $J(G)$.

The converse of Theorem 3.1 is true in some cases as shown by the following corollary.

Corollary 3.2. *Let G be either a cactus graph or a bipartite graphs or a chordal graph. Assume further that G is unmixed. Then the following are equivalent:*

- (1) $R/I(G)$ is Cohen-Macaulay;
- (2) G is vertex decomposable;
- (3) $J(G)$ is weakly polymatroidal (in some ordering of variables);
- (4) $J(G)^s$ is weakly polymatroidal (in some ordering of variables) for all $s \geq 1$.

In particular, if Δ is the independence complex of either a cactus graph or a bipartite graph or a chordal graph, then I_Δ^\vee is weakly polymatroidal in some ordering of variables if and only if Δ is vertex decomposable and pure.

Proof. (1) \Rightarrow (4) Suppose that $R/I(G)$ is Cohen-Macaulay. If G is either a cactus graph or a chordal graph, then $J(G)^s$ is weakly polymatroidal by [16, Theorem 4.3] and [17, Theorem 1.7] respectively. If G is bipartite, then G comes from a finite poset P , see [12, Theorem 9.1.13]. Using the notation in [5], we may write $I(G) = I_2(P)$. From this it follows that $(I_\Delta^\vee)^s = (H_2(P))^s$ is weakly polymatroidal for all $s \geq 1$ by [5, Theorem 2.2].

(4) \Rightarrow (3) Automatically.

(3) \Rightarrow (2) It follows from Theorem 3.1.

(2) \Rightarrow (1) Every vertex decomposable graph is sequentially Cohen-Macaulay and an unmixed sequentially Cohen-Macaulay graph is a Cohen-Macaulay graph. \square

Corollary 3.3. *Let C be a cycle of size 5. Then $\text{reg}(J(C)^{(s)}) = \text{reg}(J(C)^s) = 3s$.*

Proof. It is known that if C is a cycle of size n , then $R/I(C)$ is Cohen-Macaulay if and only if $R/I(C)$ is sequentially Cohen-Macaulay if and only if $n \in \{3, 5\}$, see [6, Proposition 4.1]. By this fact the result follows from Theorem 2.4 together with Corollary 3.2. \square

We also have the following partial converse of Theorem 3.1.

Proposition 3.4. *Let Δ be a pure simplicial complex on $[n]$. If 1 is a shedding vertex of Δ such that $I_{\Delta_i}^\vee$ is weakly polymatroidal for $i = 1, 2$, then I_Δ^\vee is weakly polymatroidal. Here, $\Delta_1 := \text{lk}_\Delta(1)$ and $\Delta_2 := \text{del}_\Delta(1)$.*

Proof. Since 1 is a shedding vertex of Δ , we have

$$G(I_\Delta^\vee) = G(I_{\Delta_1}^\vee) \sqcup \{x_1 u : u \in G(I_{\Delta_2}^\vee)\}.$$

Note that $\text{supp}(u) \subseteq \{2, \dots, n\}$ for any $u \in G(I_{\Delta_1}^\vee) \sqcup G(I_{\Delta_2}^\vee)$.

Let $\mathbf{u} = x_1^{a_1} \cdots x_n^{a_n}$ and $\mathbf{v} = x_1^{b_1} \cdots x_n^{b_n}$ be monomials belonging to $G(I_\Delta^\vee)$ satisfying $a_1 = b_1, \dots, a_{i-1} = b_{i-1}$ and $a_i < b_i$. We need to find a monomial $\mathbf{w} \in G(I_\Delta^\vee)$ such that $\mathbf{w} = x_i \mathbf{u} / x_j$ for some $j > i$. If either $\{\mathbf{u}, \mathbf{v}\} \subseteq G(I_{\Delta_1}^\vee)$ or $\{\mathbf{u}, \mathbf{v}\} \subseteq \{x_1 u : u \in G(I_{\Delta_2}^\vee)\}$, the existence of \mathbf{w} follows by the assumption that $I_{\Delta_i}^\vee$ is weakly polymatroidal for $i = 1, 2$. So we only need to consider the case that $\mathbf{u} \in G(I_{\Delta_1}^\vee)$ and $\mathbf{v} \in \{x_1 u : u \in G(I_{\Delta_2}^\vee)\}$. Note that $i = 1$ in this case. Moreover, since 1 is a shedding vertex, there exists $\mathbf{v}_1 \in G(I_{\Delta_2}^\vee)$ such that \mathbf{u} divides \mathbf{v}_1 . This implies $\mathbf{v}_1 = \mathbf{u} x_j$ for some $j > 1$ by the purity of Δ . Hence $\mathbf{w} := x_1 \mathbf{v}_1 = x_1 \mathbf{u} / x_j$ meets the requirement and so we are done. \square

Proposition 3.4 together with Corollary 3.2 leads us to present the following.

Conjecture 1. *If Δ is a vertex decomposable pure simplicial complex, then I_Δ^\vee is weakly polymatroidal.*

A weak form of this conjecture is:

Conjecture 2. *If G is an unmixed vertex decomposable graph, then $J(G)$ is weakly polymatroidal.*

The condition that G is unmixed is necessary in the above conjecture as shown by the following example.

Example 3.5. A *star graph* is a graph in which exactly one vertex has degree at least 2. Let G be a star graph with more than three vertices of degree 1. Then G is vertex decomposable, but $J(G)$ is not weakly polymatroidal in any ordering of variables.

The following results are other examples in support of Conjecture 2.

Proposition 3.6. *Let G be a Cohen-Macaulay graph of girth at least 5. Then $J(G)$ is weakly polymatroidal.*

Proof. By [15, Theorem 2.4], we have G is a \mathcal{PC} graph. Recall an induced 5-cycle of G is *basic* if it does not contain two adjacent vertices of degree three or more in G , and an edge is a *pendant* if it contains a vertex of degree 1. Let C^1, \dots, C^k be the set of all basic 5-cycles of G and let L^1, \dots, L^l be the set of all pendants of G . Recall that G is a \mathcal{PC} graph if $V(G)$ can be partitioned into $V(G) = V(C^1) \sqcup \cdots \sqcup V(C^k) \sqcup V(L^1) \sqcup \cdots \sqcup V(L^l)$.

Label the vertices of C^i successively by $x_{i1}, x_{i4}, x_{i2}, x_{i3}, x_{i5}$ such that x_{i3}, x_{i4}, x_{i5} are vertices of degree 2 for $i = 1, \dots, k$, and let y_{i1}, y_{i2} be vertices of L^i such that y_{i2} has degree 1 for $i = 1, \dots, l$. We work with the following ordering of variables:

$$x_{11} > \cdots > x_{15} > \cdots > x_{k1} > \cdots > x_{k5} > y_{11} > y_{12} > \cdots > y_{l1} > y_{l2}.$$

To prove $J(G)$ is weakly polymatroidal, we let f, g be minimal monomial generators of $J(G)$ with $f \neq g$, and let z be a variable such that $\deg_{z'} f = \deg_{z'} g$ for $z' > z$ and $\deg_z g < \deg_z f$. We need to find a variable $w < z$ such that $zg/w \in J(G)$. To this end, for each monomial $h \in J(G)$, we write h as following:

$$h = h(C^1) \cdots h(C^k)h(L^1) \cdots h(L^l),$$

where $h(C^i)$ and $h(L^j)$ are monomials such that $\text{supp}(h(C^i)) \subseteq V(C^i)$ for $i = 1, \dots, k$ and $\text{supp}(h(L^j)) \subseteq V(L^j)$ for $j = 1, \dots, l$. We consider the following cases:

Case 1 $z = x_{i1}$ for some i : Then $g(C^i) \in \{x_{i2}x_{i4}x_{i5}, x_{i3}x_{i4}x_{i5}\}$. We set $w = x_{i4}$ if $g(C^i) = x_{i2}x_{i4}x_{i5}$ and set $w = x_{i5}$ if $g(C^i) = x_{i3}x_{i4}x_{i5}$. Then $zg/w \in J(G)$.

Case 2 $z = x_{i2}$ for some i : Then $g(C^i) \in \{x_{i1}x_{i4}x_{i3}, x_{i3}x_{i4}x_{i5}\}$. We set $w = x_{i4}$ if $g(C^i) = x_{i1}x_{i4}x_{i3}$, and set $w = x_{i3}$ and set $w = x_{i4}$ if $g(C^i) = x_{i3}x_{i4}x_{i5}$. Then $zg/w \in J(G)$.

Case 3 $z = x_{i3}$ for some i : Then $g(C^i) = x_{i1}x_{i5}x_{i2}$. We set $w = x_{i5}$.

Case 4 $z = x_{i4}$ for some i : This case is impossible.

Case 5 $z = x_{i5}$ for some i : This case is impossible again.

Case 6 $z = y_{i1}$ for some i : Then $h(L_i) = y_{i2}$. We set $w = y_{i2}$.

Case 7 $z = y_{i2}$ for some i : This case is impossible again.

Thus, in all possible cases, we find a variable w which meets the requirement. This completes the proof. \square

A graph G is called *very well-covered* if $|V(G)|$ is even and every minimal vertex cover of G contains exactly $\frac{|V(G)|}{2}$ vertices. Clearly, every very well-covered graph is unmixed, i.e. well-covered.

Proposition 3.7. *Let G be a Cohen-Macaulay very well-covered graph. Then $J(G)$ is weakly polymatroidal.*

Proof. In view of [18, Lemma 3.1], there is a relabeling of vertices

$$V(G) = \{x_1, \dots, x_n, y_1, \dots, y_n\}$$

such that the following five conditions hold:

(i) $X = \{x_1, \dots, x_n\}$ is a minimal vertex cover of G and $Y = \{y_1, \dots, y_n\}$ is an independent set of G ;

(ii) $\{x_i, y_i\} \in E(G)$ for $i = 1, \dots, n$;

(iii) if $\{z_i, x_j\}, \{y_j, x_k\} \in E(G)$, then $\{z_i, x_k\} \in E(G)$ for distinct i, j, k and for $z_i \in \{x_i, y_i\}$;

(iv) if $\{x_i, y_j\} \in E(G)$, then $\{x_i, x_j\} \notin E(G)$;

(v) if $\{x_i, y_j\} \in E(G)$, then $i \leq j$.

We will show that $J(G)$ is weakly polymatroidal in the ordering: $x_1 > x_2 > \dots > x_n > y_1 > y_2 > \dots > y_n$. Let f, g be minimal monomial generators of $J(G)$ with $f \neq g$. Then we may write f and g as follows:

$$f = \prod_{z \in C} z = x_1^{a_1} \cdots x_n^{a_n} y_1^{a_{n+1}} \cdots y_n^{a_{n+n}}$$

and

$$g = \prod_{z \in D} z = x_1^{b_1} \cdots x_n^{b_n} y_1^{b_{n+1}} \cdots y_n^{b_{n+n}}.$$

Here, C and D are minimal vertex covers of G , and $a_i, b_i \in \{0, 1\}$ for all i . Note that $a_i + a_{n+i} = 1$ and $b_i + b_{n+i} = 1$ for $i = 1, \dots, n$, there exists $1 \leq k \leq n$ such that $a_i = b_i$ for $i = 1, \dots, k-1$ and $a_k > b_k$. This actually means that $x_k \in C$ and $x_k \notin D$. From this it follows that $y_k \in D$. We claim that $A := (D \cup \{x_k\}) \setminus \{y_k\}$ is also a vertex cover of G .

Let $e = \{z_1, z_2\} \in E(G)$. If $y_k \notin e$, then it is clear that $A \cap e \neq \emptyset$. So we assume that $e = \{x_j, y_k\}$. By (v), we have $j \leq k$. If $j = k$, then $A \cap e = \{x_k\}$. So we may assume further that $j < k$. Since $y_k \notin C$, it follows that $x_j \in C$ and so $b_j = a_j = 1$. This implies $x_j \in D$ and so $A \cap e \neq \emptyset$. Thus, we prove the claim and it follows that $x_k g / y_k = \prod_{z \in A} z$ is a minimal monomial generator of $J(G)$, completing the proof. \square

Let G be a graph and $s \geq 1$ an integer. In [22], the graph G_s is constructed so that $J(G_s)$ is the polarization of $J(G)^{(s)}$.

Corollary 3.8. *Let G be a Cohen-Macaulay very well-covered graph. Then $J(G)^{(s)}$ is weakly polymatroidal for all $s \geq 1$.*

Proof. It is immediate from the definitions that if I is a monomial ideal generated in one degree then I is weakly polymatroidal if and only if the polarization of I is weakly polymatroidal. By [22, Proposition 3.1], if G is a Cohen-Macaulay very well-covered graph then so is G_s . Now, the result follows from Proposition 3.7. \square

4. POWERS OF COVER IDEALS OF CLIQUE-WHISKERING GRAPHS

Let $W = G^\pi$ be the fully clique-whiskering of some graph G by some clique vertex-partition π of G . In this section we prove $J(W)^s$ is weakly polymatroidal for all $s \geq 1$. As a consequence, we have $\text{reg}(J(W)^s) = \text{reg}(J(W)^{(s)}) = s|V(G)|$ for all $s \geq 1$.

For convenience, we introduce the notions of a simplicial co-complex and the face ideal of a simplicial co-complex with respect to a partition.

Definition 4.1. Let V be a finite set. We say that a collection ∇ of subsets of V is a *simplicial co-complex* on V if whenever $F \in \nabla$ and $F \subseteq G \subseteq V$ one has $G \in \nabla$. Assume that V has a partition $V = V_1 \sqcup \dots \sqcup V_t$ and assume that $V_i = \{x_{i1}, \dots, x_{ik_i}\}$ for $i = 1, \dots, t$. For each face $F \in \nabla$, we put

$$u_F := \prod_{x_{ij} \in F} x_{ij} \prod_{i \in [t]} y_i^{k_i - |F \cap V_i|}.$$

Then the face ideal of ∇ with respect to this partition is the following monomial ideal J in the polynomial ring $k[x_{11}, \dots, x_{1k_1}, \dots, x_{t1}, \dots, x_{tk_t}, y_1, \dots, y_t]$:

$$J = (u_F : F \in \nabla).$$

Proposition 4.2. *Let ∇ be a simplicial co-complex on V and assume that V has a partition $V = V_1 \sqcup \dots \sqcup V_t$ with $V_i = \{x_{i1}, \dots, x_{ik_i}\}$ for $i = 1, \dots, t$. Denote by J the face ideal of ∇ with respect to this partition. Then J^s is weakly polymatroidal for each $s \geq 1$ in the ordering: $x_{11} > \dots > x_{1k_1} > \dots > x_{t1} > \dots > x_{tk_t} > y_1 > \dots > y_t$.*

Proof. Let α, β be the minimal monomial generators of J^s with $\alpha \neq \beta$. We may write

$$\alpha = u_{F_1} u_{F_2} \cdots u_{F_s} = u_1 \cdots u_t y_1^{sk_1 - \deg(u_1)} \cdots y_t^{sk_t - \deg(u_t)},$$

and

$$\beta = u_{G_1} u_{G_2} \cdots u_{G_s} = v_1 \cdots v_t y_1^{sk_1 - \deg(v_1)} \cdots y_t^{sk_t - \deg(v_t)}.$$

Here, $F_i \in \nabla, G_i \in \nabla$ for all $i \in [s]$, and $\text{supp}(u_i) \cup \text{supp}(v_i) \subseteq V_i$ for $i = 1, \dots, t$. Since $\alpha \neq \beta$, there exists a variable $z < y_1$ such that $\deg_w(\alpha) = \deg_w(\beta)$ for all $w < z$ and $\deg_z(\alpha) < \deg_z(\beta)$. Suppose that $z = x_{ij}$, where $1 \leq i \leq t$ and $1 \leq j \leq k_i$. Then $\deg_{x_{ij}} \alpha \leq s - 1$ and so there exists $l \in [s]$, such that $x_{ij} \notin F_l \cap V_i$. Say $l = 1$. Put $F'_1 := F_1 \cup \{x_{ij}\}$ and let

$$\gamma := u_{F'_1} u_{F_2} \cdots u_{F_s}.$$

Then $\gamma = x_{ij}\alpha/y_i$ and $\gamma \in J^s$. From this it follows that J^s is weakly polymatroidal. \square

Theorem 4.3. *Let $W = G^\pi$, where G is a simple graph and $\pi = (W_1, \dots, W_t)$ is a clique vertex-partition of G . Then $J(W)^s$ is weakly polymatroidal for every $s \geq 1$.*

Proof. Let $\nabla := \{C \cap V(G) : C \text{ is a minimal vertex cover of } W\}$. Then ∇ is a simplicial co-complex on $V(G)$. Moreover, $J(W)$ coincides with the face ideal of ∇ with respect to the partition π . Now, the result follows from Proposition 4.2. \square

Theorem 4.3 together with Theorem 3.1 implies G^π is vertex decomposable. This recovers [3, Theorem 3.3]. Other consequences of Theorem 4.3 are as follows:

Corollary 4.4. *Let $W = G^\pi$ be as in Theorem 3.1. Then $\text{reg}(J(W)^{(s)}) = \text{reg}(J(W)^s) = s|V(G)|$ for all $s \geq 1$.*

Proof. It follows from Theorem 4.3 that $\text{reg}(J(W)^s) = s|V(G)|$. By [21, Corollary 4.4], we have $\text{reg}(J(W)^{(s)}) = s|V(G)|$. \square

Corollary 4.5. *Let W be the whisker graph of some graph. Then both $J(W)^{(s)}$ and $J(W)^s$ are weakly polymatroidal for all $s \geq 1$.*

Proof. We have that $J(W)^s$ is weakly polymatroidal by Theorem 4.3 and that $J(W)^{(s)}$ is weakly polymatroidal by Corollary 3.8. \square

Acknowledgment: This research is supported by NSFC (No. 11971338). We would like to express our sincere thanks to the referee, who reads the paper carefully and helps us kill many minor errors of the original paper.

REFERENCES

- [1] A. Banerjee, S. Beyarslan, H. T. Hà: *Regularity of edge ideals and their Powers*, Springer Proceedings in Mathematics and Statistics, **277** (2019), 17–52.
- [2] A. Björner, M. Wachs: *Shellable nonpure complexes and posets. I*, Trans. Amer. Math. Soc. **348** (1996), 1299–1327.
- [3] D. Cook II, U. Nagel: *Cohen-Macaulay graphs and face vectors of flag complexes*, SIAM J. Discrete Math. **26** (2012), 89–101.

- [4] L. X. Dung, T. T. Hien, H. D. Nguyen, T. N. Trung: *Regularity and Koszul property of symbolic powers of monomial ideals*, Math. Z. **298** (2021), 1487–1522.
- [5] V. Ene, J. Herzog, F. Mohammadi: *Monomial ideals and toric rings of Hibi type arising from a finite poset*, European Journal of Combinatorics, **2011** (32), 404–421.
- [6] A. C. Francisco; A. Van. Tuyl: *Sequentially Cohen-Macaulay edge ideals*, Proc. Amer. Math. Soc. **135** (2007), 2327–2337.
- [7] Y. Gu, H. T. Hà, J.W. Skelton: *Symbolic powers of cover ideals of graphs and Koszul property*, Internat. J. Algebra Comput. **31** (2021), 865–881.
- [8] J. Guo, M. Li, T. Wu: *A new view toward vertex decomposable graphs*, Discrete Mathematics **345** (2022), 112953
- [9] A. V. Jayanthan, R. Kumar: *Regularity of symbolic powers of edge ideals*, Journal of Pure and Applied Algebra **224** (2020), 106306
- [10] M. Kokubo, T. Hibi: *Weakly polymatroidal ideals*, Algebra Colloq. **13** (2006), 711–720.
- [11] A. Kumar, R. Kumar, R. Sarkar, S. Selvaraja: *Symbolic powers of certain cover ideals of graphs*, Acta Mathematica Vietnamica **46** (2021), 599–611
- [12] J. Herzog, T. Hibi: *Monomial Ideals*, Graduate Texts in Mathematics 260, Springer, London-Dordrecht-Heidelberg-New York, 2011.
- [13] J. Herzog, T. Hibi, N. V. Trung: *Symbolic powers of monomial ideals and vertex cover algebras*, Adv. Math. **210** (2007), 304–322.
- [14] T. Hibi, A. Higashitani, K. Kimura, A.B. O’Keefe: *Algebraic study on Cameron-Walker graphs*, J. Algebra **422** (2015), 257–269.
- [15] D.T. Hoang, N.C. Minh, T.N. Trung: *Cohen-Macaulay graphs with large girth*, J. Algebra Appl. **14** (2015), 1550112
- [16] F. Mohammadi: *Powers of the vertex cover ideals*, Collect. Math. **65** (2014), 169–181.
- [17] F. Mohammadi: *Powers of the vertex cover ideal of a chordal graph*, Comm. Algebra **39** (2011), 3753–3764.
- [18] M. Mahmoudi, A. Mousivand, M. Crupi, G. Rinaldo, N. Terai, S. Yassemi: *Vertex decomposability and regularity of very well-covered graphs*, J. Pure Appl. Algebra **215** (2011), 2473–2480.
- [19] R. Nanduri: *On regularity of symbolic Rees algebras and symbolic powers of vertex cover ideals of graphs*, Proc. Amer. Math. Soc. **150** (2022), 1955–1965.
- [20] J. Provan, L. Billera: *Decompositions of simplicial complexes related to diameters of convex polyhedra*, Math. Oper. Res. **5** (1980), 576–594.
- [21] S. Selvaraja: *Symbolic powers of vertex cover ideals*, Internat. J. Algebra Comput. **30** (2020), 1167–1183.
- [22] S. A. Seyed Fakhari: *Symbolic powers of cover ideal of very well-covered and bipartite graphs*, Proc. Amer. Math. Soc. **146** (2018), 97–110
- [23] S. A. Seyed Fakhari: *Regularity of symbolic powers of cover ideals of graphs*. Collect. Math. **70** (2019), 187–195.
- [24] S. A. Seyed Fakhari: *On the minimal free resolution of symbolic powers of cover ideals of graphs*, Proc. Amer. Math. Soc. **149** (2021), 3687–3698.
- [25] N. V. Trung, T. M. Tuan: *Equality of ordinary and symbolic powers of Stanley-Reisner ideals*, J. Algebra **328** (2011), 77–93.

DANCHENG LU, SCHOOL OF MATHEMATICAL SCIENCE, SOOCHOW UNIVERSITY, P.R.CHINA
E-mail address: ludancheng@suda.edu.cn

ZEXIN WANG, SCHOOL OF MATHEMATICAL SCIENCE, SOOCHOW UNIVERSITY, P.R.CHINA
E-mail address: zexinwang6@outlook.com