

Concavity of spacetimes

Tobias Beran¹ Darius Erös¹ Shin-ichi Ohta² Felix Rott³

September 30, 2025

Abstract Motivated by recent breathtaking progress in the synthetic study of Lorentzian geometry, we investigate the local concavity of time separation functions on Finsler spacetimes as a Lorentzian counterpart to Busemann’s convexity in metric geometry. We show that a Berwald spacetime is locally concave if and only if its flag curvature is nonnegative in timelike directions. We also give another characterization of nonnegative flag curvature by the convexity of future (or past) capsules, inspired by Kristály–Kozma’s result in the positive definite case. These characterizations are new even for Lorentzian manifolds.

Keywords: Finsler spacetimes, flag curvature, concavity, convex capsules

MSC (2020): 53B30, 53C50, 53C60

1 Introduction

The aim of this article is to develop the synthetic study of Lorentzian geometry. Synthetic geometry is concerned with geometric spaces without differentiable structures, such as metric (measure) spaces. For instance, for a complete Riemannian manifold, every small triangle drawn with geodesics is thinner than its comparison triangle (with the same sidelengths) in \mathbf{R}^2 if and only if the sectional curvature is nonpositive everywhere (Alexandrov’s triangle comparison theorem; see, e.g., [BH, Theorem II.1A.6]). Then, the triangle comparison property can be adopted as a synthetic notion of nonpositive curvature; a (geodesic) metric space (X, d) is called a *CAT(0)-space* if triangles in X are thinner than \mathbf{R}^2 . In the same manner, one can define metric spaces of sectional curvature bounded below (Alexandrov spaces; see [AKP, BBI]), and metric measure spaces of Ricci curvature $\geq K$ and dimension $\leq N$ (MCP(K, N)-spaces (MCP stands for measure contraction property) and CD(K, N)-spaces (CD stands for curvature-dimension condition); see [LV, Oh1, St1, St2]). These synthetic approaches enable us to understand the influence of curvature in a geometric and intuitive way, and to investigate metric (measure) spaces arising as limit spaces of Riemannian manifolds, leading on to fruitful applications in Riemannian geometry and beyond.

Recently, there is a growing interest in synthetic approaches to Lorentzian geometry, partly motivated by the appearance of less regular spacetimes in general relativity (as solutions to the Einstein equation). For time-oriented Lorentzian manifolds (spacetimes), suitable

¹Department of Mathematics, University of Vienna, Oskar-Morgenstern-Platz 1, 1090 Vienna, Austria (tobias.beran@univie.ac.at, darius.eroes@univie.ac.at)

²Department of Mathematics, University of Osaka, Osaka 560-0043, Japan (s.ohta@math.sci.osaka-u.ac.jp); RIKEN Center for Advanced Intelligence Project (AIP), 1-4-1 Nihonbashi, Tokyo 103-0027, Japan

³SISSA, Via Bonomea 265, 34136 Trieste, Italy (frott@sissa.it)

counterparts to the triangle comparison properties are known to characterize lower and upper sectional curvature bounds in timelike directions (see [AB, BKOR, Ha]). Then, as a platform for synthetic Lorentzian geometry, Lorentzian (pre-)length spaces are introduced [KS], and the structure of Lorentzian (pre-)length spaces of curvature bounded below or above in the sense of triangle comparison is the subject of current active research (see, e.g., [BORS, BS]). We also refer to [CM, Mc, MS] for a synthetic approach to lower Ricci curvature bounds (timelike curvature-dimension condition $\text{TCD}(K, N)$), to name a few.

For metric spaces, there is also a weaker notion of nonpositive curvature than $\text{CAT}(0)$ -spaces, going back to a seminal work of Busemann [Bu], called the *convexity* or the *Busemann nonpositive curvature*. Let (X, d) be a *geodesic* space, i.e., any pair $x, y \in X$ of points can be joined by a curve $\gamma: [0, 1] \rightarrow X$ such that $\gamma(0) = x$, $\gamma(1) = y$ and $d(\gamma(s), \gamma(t)) = |t - s|d(x, y)$ for all $s, t \in [0, 1]$ (then γ is called a *minimizing geodesic*). We say that (X, d) is *locally convex* if every point $x \in X$ has a neighborhood U such that, for any minimizing geodesics $\gamma_1, \gamma_2: [0, 1] \rightarrow X$ included in U , the function $t \mapsto d(\gamma_1(t), \gamma_2(t))$ is convex. If we can take $U = X$, (X, d) is said to be *globally convex*. For Riemannian manifolds, local convexity is equivalent to having the nonpositive sectional curvature everywhere.

An important feature of Busemann's convexity is that it does not rule out Finsler manifolds. For example, strictly convex Banach spaces are clearly globally convex, while only Hilbert spaces are $\text{CAT}(0)$ among Banach spaces. Therefore, it had been an intriguing problem to characterize convex spaces among Finsler manifolds. It was first shown in [KVK] that a Berwald manifold, a special kind of Finsler manifold, of nonpositive flag curvature is locally convex (flag curvature in Finsler geometry corresponds to sectional curvature in Riemannian geometry). Then, [KK] proved that a Berwald manifold is locally convex only if its flag curvature is nonpositive everywhere. Finally, it was established in [IL] that a Finsler manifold is locally convex if and only if it is a Berwald manifold of nonpositive flag curvature. We refer to [FG, HY] for recent studies of the structure of convex metric spaces, where we need to deal with the Finsler (non-Riemannian) nature of those spaces.

As a Lorentzian counterpart to Busemann's convexity, the concavity of time separation functions on Lorentzian pre-length spaces of nonnegative curvature in the sense of triangle comparison was proved in [BKR, Section 6]. Moreover, a globalization result for such a concavity was established in [EG] (see Remark 3.9). Then, in comparison with convexity, it is natural to ask an analog to the above characterization of convex Finsler or Berwald manifolds. Our main theorem provides an answer to this question in the setting of Berwald spacetimes.

Theorem 1.1 (Characterizations of concavity). *Let (M, L) be a Berwald spacetime. Then, the following are equivalent.*

- (I) (M, L) has nonnegative flag curvature $\mathbf{K} \geq 0$ in timelike directions, namely $\mathbf{K}(v, w) \geq 0$ for every pair of linearly independent vectors $v, w \in T_x M$ such that v is future-directed timelike.
- (II) (M, L) is locally concave.
- (III) (M, L) is locally timelike concave.
- (IV) (M, L) has convex future capsules.
- (V) (M, L) has convex past capsules.

See Definition 3.1 for the precise definition of local (timelike) concavity. Having convex future or past capsules is a condition inspired by [KK]; see Definition 4.1. Finsler spacetimes generalize usual (Lorentzian) spacetimes in the same way that Finsler manifolds generalize Riemannian manifolds, and the Berwald condition is defined in the same way as well. Finsler and Berwald spacetimes have been studied from both geometric and physical viewpoints (see Section 2).

Theorem 1.1 is new even in the case of Lorentzian manifolds. Then, the equivalence between local (timelike) concavity and nonnegative sectional curvature can be regarded as a direct generalization of the aforementioned result of Busemann [Bu].

Corollary 1.2 (Lorentzian case). *Let (M, g) be a Lorentzian spacetime. Then, the following are equivalent.*

- (I) (M, g) has nonnegative sectional curvature in timelike directions.
- (II) (M, g) is locally concave.
- (III) (M, g) is locally timelike concave.
- (IV) (M, g) has convex future capsules.
- (V) (M, g) has convex past capsules.

Our proof of Theorem 1.1 follows the lines of [KK, KVK]. Due to a common difficulty in Lorentzian geometry, the noncompactness of the indicatrix in the tangent spaces, we could not generalize the argument in [IL]. Thus, it remains an open question whether locally (timelike) concave Finsler spacetimes necessarily satisfy the Berwald condition (see Subsection 5.1 for more details).

This article is organized as follows. After reviewing the basics of Finsler spacetimes in Section 2, we prove the equivalence between concavity and nonnegative flag curvature in Section 3. We then give another characterization of nonnegative flag curvature by convex capsules in Section 4. Finally, Section 5 is devoted to discussing some open problems on concavity, compared to the convexity of metric spaces.

2 Finsler spacetimes

We review the basics of Finsler spacetimes (see [BEE, ON] for the standard theory of Lorentzian spacetimes). Our definition of Finsler spacetimes follows Beem's one [Be] (see [LMO1, §2.1] for the relationship with other definitions). Concerning recent development, we refer to [LMO1, LMO2] for singularity and comparison theorems, [COO, LMO3] for timelike splitting theorems, and to [BO] for the timelike curvature-dimension condition.

2.1 Lorentz–Finsler manifolds

Throughout the article, let M be a connected C^∞ -manifold without boundary of dimension $n \geq 2$.

Definition 2.1 (Lorentz–Finsler structures). A *Lorentz–Finsler structure* on M is a function $L: TM \rightarrow \mathbf{R}$ satisfying the following conditions:

- (1) $L \in C^\infty(TM \setminus \{0\})$;
- (2) $L(cv) = c^2 L(v)$ for all $v \in TM$ and $c > 0$;
- (3) For any $v \in TM \setminus \{0\}$, the symmetric matrix

$$(g_{ij}(v))_{i,j=1}^n := \left(\frac{\partial^2 L}{\partial v^i \partial v^j}(v) \right)_{i,j=1}^n \quad (2.1)$$

is non-degenerate with signature $(-, +, \dots, +)$.

Then, we call (M, L) a $(C^\infty\text{-})$ Lorentz-Finsler manifold.

For each $v \in T_x M \setminus \{0\}$, we can define a Lorentzian metric g_v on $T_x M$ via (2.1) as

$$g_v \left(\sum_{i=1}^n a_i \frac{\partial}{\partial x^i} \Big|_x, \sum_{j=1}^n b_j \frac{\partial}{\partial x^j} \Big|_x \right) := \sum_{i,j=1}^n g_{ij}(v) a_i b_j.$$

Note that we have $g_v(v, v) = 2L(v)$ by Euler's homogeneous function theorem.

A tangent vector $v \in TM$ is said to be *timelike* (resp. *null*) if $L(v) < 0$ (resp. $L(v) = 0$). We say that v is *lightlike* if it is null and nonzero, and *causal* if it is timelike or lightlike. *Spacelike* vectors are $v \in TM$ such that $L(v) > 0$ or $v = 0$. Denote by $\Omega'_x \subset T_x M$ the set of timelike vectors. For causal vectors v , we define

$$F(v) := \sqrt{-2L(v)} = \sqrt{-g_v(v, v)}. \quad (2.2)$$

Definition 2.2 (Finsler spacetimes). If a Lorentz-Finsler manifold (M, L) admits a smooth timelike vector field X , then (M, L) is said to be *time-oriented* (by X). A time-oriented Lorentz-Finsler manifold is called a *Finsler spacetime*.

In a Finsler spacetime time-oriented by X , a causal vector $v \in T_x M$ is said to be *future-directed* if it lies in the same connected component of $\overline{\Omega'_x} \setminus \{0\}$ as $X(x)$. We denote by $\Omega_x \subset \Omega'_x$ the set of future-directed timelike vectors, and define

$$\Omega := \bigcup_{x \in M} \Omega_x.$$

A C^1 -curve in M is said to be *timelike* (resp. *causal*) if its tangent vector is always timelike (resp. causal). Henceforth, unless explicitly stated otherwise, all causal curves are assumed to be future-directed (i.e., its tangent vector is future-directed causal).

Next, we introduce the covariant derivative. Define

$$\gamma_{jk}^i(v) := \frac{1}{2} \sum_{l=1}^n g^{il}(v) \left(\frac{\partial g_{lk}}{\partial x^j} + \frac{\partial g_{jl}}{\partial x^k} - \frac{\partial g_{jk}}{\partial x^l} \right)(v)$$

for $i, j, k = 1, \dots, n$ and $v \in TM \setminus \{0\}$, where $(g^{ij}(v))$ is the inverse matrix of $(g_{ij}(v))$;

$$G^i(v) := \sum_{j,k=1}^n \gamma_{jk}^i(v) v^j v^k, \quad N_j^i(v) := \frac{1}{2} \frac{\partial G^i}{\partial v^j}(v)$$

for $v \in TM \setminus \{0\}$, and $G^i(0) = N_j^i(0) := 0$. Moreover, we set

$$\Gamma_{jk}^i(v) := \gamma_{jk}^i(v) - \frac{1}{2} \sum_{l,m=1}^n g^{il}(v) \left(\frac{\partial g_{lk}}{\partial v^m} N_j^m + \frac{\partial g_{jl}}{\partial v^m} N_k^m - \frac{\partial g_{jk}}{\partial v^m} N_l^m \right)(v) \quad (2.3)$$

on $TM \setminus \{0\}$, and the *covariant derivative* of a vector field $Y = \sum_{i=1}^n Y^i(\partial/\partial x^i)$ is defined as

$$D_v^w Y := \sum_{i,j=1}^n \left\{ v^j \frac{\partial Y^i}{\partial x^j}(x) + \sum_{k=1}^n \Gamma_{jk}^i(w) v^j Y^k(x) \right\} \frac{\partial}{\partial x^i} \Big|_x$$

for $v \in T_x M$ with a *reference vector* $w \in T_x M \setminus \{0\}$. We remark that the functions Γ_{jk}^i in (2.3) are the coefficients of the *Chern(-Rund) connection*.

In the Lorentzian case, g_{ij} is constant in each tangent space (thus, $\Gamma_{jk}^i = \gamma_{jk}^i$) and the covariant derivative does not depend on the choice of a reference vector. In the Lorentz-Finsler setting, the following class is worth considering.

Definition 2.3 (Berwald spacetimes). A Finsler spacetime (M, L) is said to be of *Berwald type* (or called a *Berwald spacetime*) if Γ_{jk}^i is constant on the slit tangent space $T_x M \setminus \{0\}$ for every x .

By definition, the covariant derivative on a Berwald spacetime is defined independently of the choice of a reference vector. Thus, in the sequel, reference vectors will be omitted in Berwald spacetimes. An important property of Berwald spacetimes is that, for any C^1 -curve $\eta: [0, 1] \rightarrow M$ whose velocity does not vanish, the parallel transport along η gives a linear isometry between $(T_{\eta(0)} M, L)$ and $(T_{\eta(1)} M, L)$, i.e., for any vector field V along η such that $D_{\dot{\eta}} V \equiv 0$, $L(V)$ is constant (see, e.g., [Oh2, Proposition 6.5] in the positive definite case). In particular, all tangent spaces are mutually linearly isometric.

Remark 2.4 (Metrizability). In the positive definite case, Szabó showed that a Finsler manifold of Berwald type (M, F) admits a Riemannian metric h whose Levi-Civita connection coincides with the Chern connection of F , i.e., the Christoffel symbols of h coincide with Γ_{jk}^i of F (see [Sz], [BCS, Exercise 10.1.4]). This is called the (*Riemannian*) *metrizability theorem*. It is not known whether the metrizability can be generalized to Berwald spacetimes. In [FHPV], some counter-examples were constructed for Lorentz-Finsler structures defined only on a subset of TM . Their discussion is not applicable to Lorentz-Finsler structures defined on the entire tangent bundle as in Definition 2.1.

The *geodesic equation* is $D_{\dot{\eta}}^{\dot{\eta}} \dot{\eta} \equiv 0$. Then, the *exponential map* is defined in the same way as the Riemannian (or Finsler) case. For C^1 -vector fields V, W along a nonconstant geodesic η , we have

$$\frac{d}{dt} [g_{\dot{\eta}}(V, W)] = g_{\dot{\eta}}(D_{\dot{\eta}}^{\dot{\eta}} V, W) + g_{\dot{\eta}}(V, D_{\dot{\eta}}^{\dot{\eta}} W). \quad (2.4)$$

Moreover, for C^1 -vector fields V, W along a C^1 -curve η such that V is nowhere vanishing,

$$\frac{d}{dt} [g_V(V, W)] = g_V(D_{\dot{\eta}}^V V, W) + g_V(V, D_{\dot{\eta}}^V W) \quad (2.5)$$

(see, e.g., [LMO1, (3.1), (3.2)], [Oh2, Lemmas 4.8, 4.9]).

A C^∞ -vector field J along a geodesic η is called a *Jacobi field* if it satisfies the *Jacobi equation* $D_\eta^\eta D_\eta^\eta J + R_\eta(J) = 0$, where

$$R_v(w) := \sum_{i,j=1}^n R_j^i(v) w^j \frac{\partial}{\partial x^i} \Big|_x$$

for $v, w \in T_x M$ and

$$R_j^i(v) := \frac{\partial G^i}{\partial x^j}(v) - \sum_{k=1}^n \left\{ \frac{\partial N_j^i}{\partial x^k}(v) v^k - \frac{\partial N_j^i}{\partial v^k}(v) G^k(v) + N_k^i(v) N_j^k(v) \right\}$$

is the *curvature tensor*. A Jacobi field is also characterized as the variational vector field of a geodesic variation. One can further expand this as $R_j^i(v) = \sum_{k,l=1}^n R_{ljk}^i(v) v^k v^l$, where

$$R_{ljk}^i := \left(\frac{\partial \Gamma_{kl}^i}{\partial x^j} - \sum_{m=1}^n N_j^m \frac{\partial \Gamma_{kl}^i}{\partial v^m} \right) - \left(\frac{\partial \Gamma_{jl}^i}{\partial x^k} - \sum_{m=1}^n N_k^m \frac{\partial \Gamma_{jl}^i}{\partial v^m} \right) + \sum_{m=1}^n (\Gamma_{jm}^i \Gamma_{kl}^m - \Gamma_{km}^i \Gamma_{jl}^m)$$

(see [BCS, (3.3.2)], [Oh2, (5.10)] in the positive definite setting).

For $v \in \Omega_x$ and $w \in T_x M$ linearly independent of v , define the *flag curvature* of the 2-plane (*flag*) $v \wedge w$ with *flagpole* v as

$$\mathbf{K}(v, w) := \frac{g_v(R_v(w), w)}{g_v(v, v)g_v(w, w) - g_v(v, w)^2}.$$

Note that the denominator is negative (by the reverse Cauchy–Schwarz inequality for g_v as in (3.3) below). We say that the *timelike flag curvature* is nonnegative or $\mathbf{K} \geq 0$ holds in *timelike directions* if $\mathbf{K}(v, w) \geq 0$ for all v and w as above.

Remark 2.5 (Sign of \mathbf{K}). The sign of \mathbf{K} (in timelike directions) above is the same as [BEE, BKOR] and opposite to [LMO1]. Thus, the nonnegative flag curvature $\mathbf{K} \geq 0$ in timelike directions corresponds to the nonpositive timelike curvature in the sense of [AB, BKR, KS] (cf. [KS, Example 4.9], [BKOR, Remark 2.2]). Note also that $\mathbf{K} \geq 0$ holds in timelike directions for the de Sitter space (Lorentzian pseudo-sphere) and product spacetimes $(\mathbf{R} \times N, -dt^2 + h)$ where (N, h) is an Hadamard manifold.

2.2 Causality conditions and geodesics

Given $x, y \in M$, we write $x \ll y$ (resp. $x < y$) if there is a timelike (resp. causal) curve from x to y , and $x \leq y$ means that $x = y$ or $x < y$. Then, we define the *chronological past* and *future* of x by

$$I^-(x) := \{y \in M \mid y \ll x\}, \quad I^+(x) := \{y \in M \mid x \ll y\},$$

and the *causal past* and *future* of x by

$$J^-(x) := \{y \in M \mid y \leq x\}, \quad J^+(x) := \{y \in M \mid x \leq y\}.$$

Definition 2.6 (Causality conditions). Let (M, L) be a Finsler spacetime.

(1) (M, L) is said to be *chronological* if $x \notin I^+(x)$ for all $x \in M$.

- (2) We say that (M, L) is *causal* if there is no closed causal curve.
- (3) (M, L) is said to be *strongly causal* if, for all $x \in M$, every neighborhood U of x contains another neighborhood V of x such that no causal curve intersects V more than once. In particular, the topology contains a basis of causally convex sets.
- (4) We say that (M, L) is *globally hyperbolic* if it is strongly causal and, for any $x, y \in M$, $J^+(x) \cap J^-(y)$ is compact.

Define the *time separation* (also called the *Lorentz–Finsler distance*) $\tau(x, y)$ for $x, y \in M$ by

$$\tau(x, y) := \sup_{\eta} \mathbf{L}(\eta), \quad \mathbf{L}(\eta) := \int_0^1 F(\dot{\eta}(t)) dt,$$

where the supremum is taken over all (piecewise C^1) causal curves $\eta: [0, 1] \rightarrow M$ from x to y . We set $\tau(x, y) := 0$ if $x \not\leq y$. A curve η attaining the above supremum is said to be *maximizing*. In general, τ is only lower semi-continuous and can be infinite (see, e.g., [Mi2, Proposition 6.7]). In globally hyperbolic Finsler spacetimes, τ is finite and continuous, and any pair of points $x, y \in M$ with $x < y$ admits a maximizing geodesic from x to y (Avez–Seifert-type theorem; see [Mi2, Propositions 6.8, 6.9]).

3 Concavity of Berwald spacetimes

Let (M, L) be a Finsler spacetime. For each $x \in M$, there exists a *convex normal neighborhood* $U \subset M$ of x in the sense that, for every $y \in U$, the exponential map \exp_y gives a C^1 -diffeomorphism from an open star-shaped set $\mathcal{U}_y \subset T_x M$ onto U (see [Mi1, Theorem 4]). In particular, for any $y, z \in U$, there is a unique geodesic from y to z contained in U . In this and the following sections, since we shall deal only with local structures (governed by curvature), we are concerned with causal relations and the time separation restricted to U . They coincide with the global ones if (M, L) is strongly causal (by taking U smaller if necessary).

The following is a Lorentzian analog to the *convexity* of metric spaces (also called the *Busemann nonpositive curvature*; recall the introduction), inspired by [BKR, EG].

Definition 3.1 (Concavity). Let (M, L) be a Finsler spacetime.

- (1) We say that (M, L) is *locally timelike concave* if every $x \in M$ admits a convex normal neighborhood U such that, for any timelike geodesics $\eta, \xi: [0, 1] \rightarrow U$ with $\eta(t) \ll_U \xi(t)$ or $\eta(t) = \xi(t)$ at $t = 0, 1$, we have

$$\tau_U(\eta(t), \xi(t)) \geq (1 - t)\tau_U(\eta(0), \xi(0)) + t\tau_U(\eta(1), \xi(1))$$

for all $t \in [0, 1]$.

- (2) We say that (M, L) is *locally concave* if every $x \in M$ admits a convex normal neighborhood U such that, for any geodesics $\eta, \xi: [0, 1] \rightarrow U$ with $\eta(t) \ll_U \xi(t)$ or $\eta(t) = \xi(t)$ at $t = 0, 1$, we have

$$\tau_U(\eta(t), \xi(t)) \geq (1 - t)\tau_U(\eta(0), \xi(0)) + t\tau_U(\eta(1), \xi(1))$$

for all $t \in [0, 1]$.

Here we respectively denote by \ll_U and τ_U the timelike relation and time separation on $(U, L|_U)$.

In the sequel, for simplicity, we will suppress U in \ll_U and τ_U .

Remark 3.2 (Causal character of geodesics). Note that η, ξ in (2) above are not necessarily causal, and that local concavity clearly implies local timelike concavity. It was proved in [BKR, Proposition 6.1] that a Lorentzian pre-length space of timelike nonpositive curvature in the sense of strict causal triangle comparison is locally timelike concave (recall Remark 2.5 for the sign convention in timelike curvature bounds). In such a synthetic setting, it is natural to consider only timelike geodesics, since we do not know how to define spacelike geodesics. Then, in [EG, Definition 2.1], local timelike concavity as in (1) above is termed local concavity. For Berwald spacetimes, we will see that local timelike concavity and local concavity are, in fact, equivalent.

To show the local concavity from $\mathbf{K} \geq 0$, we can apply a similar calculation to the positive definite case (cf. [KVK], [Oh2, Theorem 8.30]), while causality needs to be addressed. A related result in the Lorentzian case can be found in [BEE, Proposition 11.15(1)].

Proposition 3.3 (Concave norm). *Let (M, L) be a Berwald spacetime of nonnegative flag curvature $\mathbf{K} \geq 0$ in timelike directions. If $\sigma: [0, 1] \times (-\varepsilon, \varepsilon) \rightarrow U$ is a variation in a convex normal neighborhood U such that $\sigma_s(t) := \sigma(t, s)$ is a geodesic for each $s \in (-\varepsilon, \varepsilon)$ and that $V(t, s) := \partial_s \sigma(t, s)$ is timelike for all (t, s) , then, for every s , the function $t \mapsto F(V(t, s))$ is concave.*

Proof. Put $T(t, s) := \partial_t \sigma(t, s)$. Observe from (2.2) and (2.5) that

$$\frac{\partial^2 [F^2(V)]}{\partial t^2} = -\frac{\partial^2 [g_V(V, V)]}{\partial t^2} = -2g_V(V, D_T D_T V) - 2g_V(D_T V, D_T V)$$

(recall that we can omit reference vectors in Berwald spacetimes). Since $V(\cdot, s)$ is a Jacobi field by the hypothesis for σ , we find

$$g_V(V, D_T D_T V) = -g_V(V, R_T(V)) = -g_V\left(V, \sum_{i,j,k,l=1}^n R_{ljik}^i(T) V^j T^k T^l \frac{\partial}{\partial x^i}\right). \quad (3.1)$$

Note that $R_{ljik}^i(T) = R_{ljik}^i(V)$ by the Berwald condition, and that

$$\begin{aligned} g_V\left(V, \sum_{i,j,k,l=1}^n R_{ljik}^i(V) V^j T^k T^l \frac{\partial}{\partial x^i}\right) &= g_V(R_V(T), T) \\ &= \mathbf{K}(V, T) \{g_V(V, V)g_V(T, T) - g_V(V, T)^2\}, \end{aligned} \quad (3.2)$$

where the first equation follows from [Oh2, Lemma 5.15(iv)] (which is valid regardless of the signature of metrics).

Now, $\mathbf{K}(V, T) \geq 0$ by hypothesis and, choosing a local chart $(x^i)_{i=1}^n$ which is g_V -orthonormal with $\partial/\partial x^1 = V/F(V)$, we observe

$$F^2(V)g_V(T, T) + g_V(V, T)^2 = V_1^2 \left(-T_1^2 + \sum_{i=2}^n T_i^2 \right) + (V_1 T_1)^2 \geq 0. \quad (3.3)$$

Hence, we obtain

$$\begin{aligned}
\frac{\partial^2[F(V)]}{\partial t^2} &= \frac{1}{2F(V)} \frac{\partial^2[F^2(V)]}{\partial t^2} - \frac{1}{F(V)} \left(\frac{\partial[F(V)]}{\partial t} \right)^2 \\
&\leq -\frac{g_V(D_TV, D_TV)}{F(V)} - \frac{1}{F(V)} \left(\frac{g_V(V, D_TV)}{F(V)} \right)^2 \\
&= -\frac{1}{F^3(V)} \left\{ F^2(V)g_V(D_TV, D_TV) + g_V(V, D_TV)^2 \right\}.
\end{aligned} \tag{3.4}$$

Since $F^2(V)g_V(D_TV, D_TV) + g_V(V, D_TV)^2 \geq 0$ by (3.3), this implies that $\partial^2[F(V)]/\partial t^2 \leq 0$. Therefore, $F(V(t, s))$ is concave in t . \square

The following two lemmas are concerned with the causality issue.

Lemma 3.4 (Timelike homotopy). *Let U be a convex normal neighborhood in a Finsler spacetime (M, L) . Then, for any timelike curves $\alpha, \beta: [0, 1] \rightarrow U$, there exists a homotopy $h: [0, 1] \times [0, 1] \rightarrow U$ such that $h(0; s) = \alpha(s)$, $h(1; s) = \beta(s)$ and $s \mapsto h(t; s)$ is timelike for each t , and that h and $\partial_s h$ are continuous.*

Proof. Put $x = \alpha(0)$, $y = \beta(0)$ and let $\eta: [0, 1] \rightarrow U$ be the geodesic from x to y . Consider a smooth timelike vector field X that exists by time orientation and, replacing it with cX for small $c > 0$ if necessary, we assume that $\exp_{\eta(t)}(sX(\eta(t))) \in U$ for all $t, s \in [0, 1]$.

We first choose a homotopy h_1 from α to the geodesic from x to $\alpha(\varepsilon)$ for small $\varepsilon > 0$, and take $v \in \Omega_x$ with $\alpha(\varepsilon) = \exp_x(v)$. Then, set

$$h_2(t; s) := \exp_x\left(s((1-t)v + tX(x))\right), \quad h_3(t; s) := \exp_{\eta(t)}\left(sX(\eta(t))\right)$$

for $t, s \in [0, 1]$. Concatenating h_1 , h_2 and h_3 , and connecting $h_3(1; s) = \exp_y(sX(y))$ and β in the same way, we obtain a homotopy that satisfies the desired property. \square

Lemma 3.5 (Timelike variation). *Let (M, L) be a Berwald spacetime of nonnegative flag curvature $\mathbf{K} \geq 0$ in timelike directions, and $U \subset M$ be a convex normal neighborhood. Given timelike curves $\alpha, \beta: [0, 1] \rightarrow U$, let $t \mapsto \sigma(t, s)$ be the geodesic from $\alpha(s)$ to $\beta(s)$ in U . Then, $\partial_s \sigma$ is timelike and $t \mapsto F(\partial_s \sigma(t, s))$ is concave for each $s \in [0, 1]$.*

Proof. Let $h(\lambda; s)$ be a homotopy between α and β given by Lemma 3.4, i.e., $h(0; s) = \alpha(s)$, $h(1; s) = \beta(s)$ and $\partial_s h$ is always timelike. For each $\lambda \in [0, 1]$, we consider the variation $\Xi(\lambda; \cdot, \cdot): [0, 1] \times [0, 1] \rightarrow U$ such that $t \mapsto \Xi(\lambda; t, s)$ is the geodesic from $\Xi(\lambda; 0, s) := h(\lambda; s)$ to $\Xi(\lambda; 1, s) := \beta(s)$ in U . Observe that $\Xi(0; t, s) = \sigma(t, s)$ and $\Xi(1; t, s) = \beta(s)$.

Now, we set

$$\lambda_0 := \sup\{\lambda \in [0, 1] \mid \partial_s \Xi(\lambda; t, s) \text{ is not timelike for some } t, s \in [0, 1]\}.$$

Note that $\lambda_0 < 1$, since $\partial_s \Xi(1; t, s) = \dot{\beta}(s)$ is timelike and being timelike is an open condition. Then, for any $\lambda \in (\lambda_0, 1]$, since $\partial_s \Xi(\lambda; t, s)$ is timelike for all $t, s \in [0, 1]$, it follows from Proposition 3.3 that $F(\partial_s \Xi(\lambda; t, s))$ is concave in t . Taking the limit as $\lambda \rightarrow \lambda_0$, we find that $F(\partial_s \Xi(\lambda_0; t, s))$ is concave in t . Hence,

$$\begin{aligned}
F(\partial_s \Xi(\lambda_0; t, s)) &\geq (1-t)F(\partial_s \Xi(\lambda_0; 0, s)) + tF(\partial_s \Xi(\lambda_0; 1, s)) \\
&= (1-t)F(\partial_s h(\lambda_0; s)) + tF(\dot{\beta}(s)) > 0
\end{aligned}$$

for all $t, s \in [0, 1]$. This shows that $\partial_s \Xi(\lambda_0; t, s)$ is also timelike for all $t, s \in [0, 1]$.

If $\lambda_0 > 0$, then we deduce from the openness of being a timelike vector that $\partial_s \Xi(\lambda; t, s)$ is timelike for all $t, s \in [0, 1]$ and $\lambda \in (\lambda_0 - \varepsilon, 1]$ for sufficiently small $\varepsilon > 0$, contradicting the definition of λ_0 . Consequently, $\lambda_0 = 0$ and $\partial_s \Xi(\lambda; t, s)$ is timelike for all $\lambda, t, s \in [0, 1]$. Taking $\lambda = 0$ implies that $\partial_s \sigma$ is timelike, and $F(\partial_s \sigma(t, s)) = F(\partial_s \Xi(0; t, s))$ is concave in t . \square

We are ready to prove (I) \Rightarrow (II) in Theorem 1.1.

Theorem 3.6 ($\mathbf{K} \geq 0$ implies concavity). *Let (M, L) be a Berwald spacetime of nonnegative flag curvature $\mathbf{K} \geq 0$ in timelike directions. Then, (M, L) is locally concave.*

Proof. Let $U \subset M$ be a convex normal neighborhood. We first consider geodesics $\eta, \xi: [0, 1] \rightarrow U$ with $\eta(0) \ll \xi(0)$ and $\eta(1) \ll \xi(1)$. Then, take the geodesics α from $\eta(0)$ to $\xi(0)$ and β from $\eta(1)$ to $\xi(1)$, and let $\sigma: [0, 1] \times [0, 1] \rightarrow U$ be the variation such that $\sigma(0, s) = \alpha(s)$, $\sigma(1, s) = \beta(s)$, and that $t \mapsto \sigma(t, s)$ is the geodesic from $\alpha(s)$ to $\beta(s)$. Then, it follows from Lemma 3.5 that $\partial_s \sigma$ is timelike and $F(\partial_s \sigma(t, s))$ is concave in t . This implies that

$$\mathbf{L}(t) := \int_0^1 F(\partial_s \sigma(t, s)) \, ds$$

is concave, and hence

$$\tau(\eta(t), \xi(t)) \geq \mathbf{L}(t) \geq (1-t)\mathbf{L}(0) + t\mathbf{L}(1) = (1-t)\tau(\eta(0), \xi(0)) + t\tau(\eta(1), \xi(1)),$$

as desired.

The case of $\eta(0) = \xi(0)$ or $\eta(1) = \xi(1)$ is obtained as the limit, thanks to the continuity of geodesics in their endpoints (in U). \square

The converse implication (from concavity to the Berwald condition and $\mathbf{K} \geq 0$) was established in the positive definite case in [IL]; however, their argument cannot apply to the Lorentzian setting (see Subsection 5.1 for more details). Here we consider a weaker result in the spirit of [KK], putting the Berwald condition in the assumption. To this end, we need another preliminary lemma.

Lemma 3.7 (Concavity yields concave norm). *Let J be a timelike Jacobi field along a timelike geodesic $\eta: [0, \delta] \rightarrow U$ in a convex normal neighborhood $U \subset M$. If τ is timelike concave in U (in the sense of Definition 3.1), then $t \mapsto F(J(t))$ is concave. Similarly, if τ is concave in U , then $F(J)$ is concave for every timelike Jacobi field J along any geodesic η in U .*

Proof. Let $\sigma: [0, \delta] \times (-\varepsilon, \varepsilon) \rightarrow U$ be any geodesic variation associated with J , i.e., $\sigma_s := \sigma(\cdot, s)$ is a geodesic for every s , $\sigma_0 = \eta$, and $\partial_s \sigma(t, 0) = J(t)$ for all t . Note that σ_s is timelike for s close to 0. Moreover, since $J(t)$ is timelike, $t \mapsto \tau(\eta(t), \sigma_s(t))$ is concave for small $s > 0$ by timelike concavity. Then, the concavity of $t \mapsto F(J(t))$ follows from

$$\lim_{s \downarrow 0} \frac{\tau(\eta(t), \sigma_s(t))}{s} = F(J(t)).$$

The latter assertion is shown in the same way. \square

The next theorem shows (III) \Rightarrow (I) in Theorem 1.1.

Theorem 3.8 (Timelike concavity implies $\mathbf{K} \geq 0$). *Let (M, L) be a Berwald spacetime. If (M, L) is locally timelike concave, then we have $\mathbf{K} \geq 0$ in timelike directions.*

Proof. Suppose to the contrary that $g_v(R_v(w), w) > 0$ holds for some linearly independent timelike vectors $v, w \in \Omega_x$. Let $\eta: [0, \delta] \rightarrow M$ be the geodesic with $\dot{\eta}(0) = w$ and J be the Jacobi field along η with $J(0) = v$ and $D_{\dot{\eta}}J(0) = 0$. Then, we have

$$D_{\dot{\eta}}D_{\dot{\eta}}J(0) = -R_w(v)$$

and, by (2.5),

$$\frac{d}{dt} [L(J(t))]_{t=0} = \frac{1}{2} \frac{d}{dt} [g_J(J(t), J(t))]_{t=0} = g_v(v, D_{\dot{\eta}}J(0)) = 0.$$

Combining these and recalling $F = \sqrt{-2L}$, we obtain

$$\begin{aligned} \frac{d^2}{dt^2} [F(J(t))]_{t=0} &= -\frac{d}{dt} \left[\frac{1}{\sqrt{-2L(J(t))}} \cdot \frac{d}{dt} [L(J(t))] \right]_{t=0} \\ &= -\frac{1}{F(v)} \cdot \frac{d}{dt} [g_J(J(t), D_{\dot{\eta}}J(t))]_{t=0} \\ &= \frac{1}{F(v)} \cdot g_v(v, R_w(v)). \end{aligned}$$

However, as we saw in (3.1) and (3.2), the Berwald condition ensures that

$$g_v(v, R_w(v)) = g_v(R_v(w), w) > 0.$$

This contradicts Lemma 3.7, and completes the proof. \square

Remark 3.9 (Globalization). (a) In the case of metric spaces, Busemann's convexity is known to enjoy the *globalization* property (called the Cartan–Hadamard theorem): If a complete, connected metric space (X, d) is locally convex, then its universal cover \tilde{X} with its induced length metric is globally convex (see [BH, Theorem II.4.1(1)] and the references therein). In particular, if (X, d) is itself a simply connected length space, then it is globally convex.

(b) In the Lorentzian setting, an analogous statement was proved in [EG, Theorem 4.9]: If a globally hyperbolic, regular Lorentzian length space X is locally concave and future one-connected, then it is globally concave. Here, global hyperbolicity replaces the metric completeness assumption, and future one-connectedness refers to the requirement that X be timelike path-connected and any pair of future-directed timelike curves with common endpoints be homotopic through a family of timelike curves. Recall from Remark 3.2 that concavity in [EG] means timelike concavity in the sense of Definition 3.1.

4 Convex capsules

We next consider another kind of characterization of the concavity condition for Berwald spacetimes. Inspired by [KK], we introduce the following.




Figure 1: The convex past capsule associated to a geodesic γ .

Definition 4.1 (Convex capsules). A Finsler spacetime (M, L) is said to have *convex future capsules* if, for every $x \in M$, there exists a convex normal neighborhood U_x of x and $r > 0$ such that the set

$$K_{\geq r}^+(\gamma) := \bigcup_{y \in \gamma} H_{\geq r}^+(y) \cap U_x$$

is geodesically convex for any geodesic γ contained in U_x , where

$$H_{\geq r}^+(y) := \{z \in M \mid \tau(y, z) \geq r\}.$$

To have *convex past capsules* is defined in the same way with

$$K_{\geq r}^-(\gamma) := \bigcup_{y \in \gamma} H_{\geq r}^-(y) \cap U_x, \quad H_{\geq r}^-(y) := \{z \in M \mid \tau(z, y) \geq r\}.$$

(To be more precise, τ in the definitions of $H_{\geq r}^\pm(y)$ should be read as τ_{U_x} .)

In [KK, Definition 3], they considered the convexity of a neighborhood of a short geodesic in a metric space, whose shape indeed resembles a capsule. In our definition, $K_{\geq r}^\pm(\gamma)$ does not look like a capsule (see Figure 1), but we keep the name and notation to make the correspondence with [KK] transparent.

Proposition 4.2 (Concavity implies convex capsules). *If a Berwald spacetime (M, L) is locally concave, then it has convex future and past capsules.*

Proof. Given $x \in M$, let U_x be a convex normal neighborhood of x and γ be any geodesic in U_x . Take $z_1, z_2 \in K := K_{\geq r}^+(\gamma) \subset U_x$ and let $\alpha: [0, 1] \rightarrow U_x$ be the geodesic from z_1 to z_2 . Since $z_1, z_2 \in K$, we find $t_1, t_2 \in [0, 1]$ such that $\tau(\gamma(t_1), z_1) \geq r$ and $\tau(\gamma(t_2), z_2) \geq r$. If $t_1 \leq t_2$, then, we infer from the assumed concavity that, for any $s \in [0, 1]$,

$$\tau(\gamma((1-s)t_1 + st_2), \alpha(s)) \geq (1-s)\tau(\gamma(t_1), z_1) + s\tau(\gamma(t_2), z_2) \geq r.$$

This shows that $\alpha(s) \in K$. If $t_2 < t_1$, then we observe from the Berwald condition that the reversal of $\gamma|_{[t_2, t_1]}$ is still a geodesic from $\gamma(t_1)$ to $\gamma(t_2)$ and the above argument shows $\alpha(s) \in K$. This completes the proof for the future case.

We can see that concavity implies convex past capsules in the same way. \square

In the converse implication, we can roughly follow the argument in [KK] ((e) \Rightarrow (a) of Theorem 1).




Figure 2: The geodesic variation σ and a tangent hyperbola.

Proposition 4.3 (Convex capsules imply $\mathbf{K} \geq 0$). *Let (M, L) be a Berwald spacetime having convex future capsules. Then, we have $\mathbf{K} \geq 0$ in timelike directions. Similarly, if (M, L) is a Berwald spacetime having convex past capsules, then $\mathbf{K} \geq 0$ in timelike directions.*

Proof. Let $x \in M$. It is sufficient to show $\mathbf{K}(v, w) \geq 0$ for any $v \in \Omega_x$ and $w \in T_x M$ satisfying $F(v) = 1$, $g_v(v, w) = 0$ and $g_v(w, w) = 1$. For small $r > 0$, let $\eta: [0, r] \rightarrow U_x$ be the geodesic with $\dot{\eta}(0) = v$ and W be the vector field along η such that $W(0) = w$ and $D_{\dot{\eta}}W \equiv 0$. (Recall that we can drop reference vectors in Berwald spacetimes.) It follows from (2.4) or (2.5) that

$$\frac{d}{dt} [g_{\dot{\eta}}(\dot{\eta}, W)] = g_{\dot{\eta}}(D_{\dot{\eta}}\dot{\eta}, W) + g_{\dot{\eta}}(\dot{\eta}, D_{\dot{\eta}}W) = 0.$$

Hence, $g_{\dot{\eta}}(\dot{\eta}(t), W(t)) = 0$ holds for all $t \in [0, r]$.

Consider the variation $\sigma: [0, r] \times (-\varepsilon, \varepsilon) \rightarrow U_x$ such that $\sigma_t(s) := \sigma(t, s)$ is the geodesic with $\dot{\sigma}_t(0) = W(t)$, and define $T := \partial_t \sigma$ and $V := \partial_s \sigma$ (thus $V(t, 0) = W(t)$). Note that σ_r is tangent to $H_{\geq r}^+(x) \cap U_x$ at $\eta(r)$ by $g_{\dot{\eta}}(\dot{\eta}(r), W(r)) = 0$ (see Figure 2). Moreover, we deduce from $g_{\dot{\eta}}(\dot{\eta}, W) \equiv 0$ and the first variation formula for $\mathbf{L}(s) := \int_0^r F(T(t, s)) dt$ (see [BO, Lemma 2.6]) that $\mathbf{L}'(0) = 0$ and $\eta(t) \notin K$ for $t \in [0, r]$, where

$$K := K_{\geq r}^+(\sigma_0) = \bigcup_{s \in (-\varepsilon, \varepsilon)} H_{\geq r}^+(\sigma_0(s)) \cap U_x.$$

Since K is a geodesically convex set (by hypothesis) including $H_{\geq r}^+(x) \cap U_x$, we infer that σ_r is also tangent to K . This implies that, by the convexity of K , σ_r does not enter the interior of K . In particular, $\sigma_r(s)$ is not in the interior of $H_{\geq r}^+(\sigma_0(s)) \cap U_x$. Hence,

$$\mathbf{L}(s) \leq \tau(\sigma_0(s), \sigma_r(s)) \leq r.$$

Combining this with $\mathbf{L}(0) = r$ and $\mathbf{L}'(0) = 0$, we obtain $\mathbf{L}''(0) \leq 0$.

For each $t \in [0, r]$, $s \mapsto T(t, s)$ is a Jacobi field along σ_t by construction. Thus, we find that

$$\left. \frac{\partial^2 [F^2(T)]}{\partial s^2} \right|_{s=0} = 2g_{\dot{\eta}}(\dot{\eta}, R_W(\dot{\eta})) - 2g_{\dot{\eta}}(D_W T, D_W T)$$

(recall $V(t, 0) = W(t)$). In the first term, by (3.1), (3.2) and $D_{\dot{\eta}}W \equiv 0$, we infer that

$$g_{\dot{\eta}}(\dot{\eta}, R_W(\dot{\eta})) = \mathbf{K}(\dot{\eta}, W) \{g_{\dot{\eta}}(\dot{\eta}, \dot{\eta})g_{\dot{\eta}}(W, W) - g_{\dot{\eta}}(\dot{\eta}, W)^2\} = -\mathbf{K}(\dot{\eta}, W).$$

As to the second term, we have

$$D_V T(t, 0) = D_T V(t, 0) = D_{\dot{\eta}} W(t) = 0.$$

Hence, by a similar calculation to (3.4) (using $F(\dot{\eta}) \equiv 1$), we obtain

$$\begin{aligned} \mathbf{L}''(0) &= \int_0^r \left\{ \frac{1}{2} \frac{\partial^2 [F^2(T)]}{\partial s^2} \Big|_{s=0} - \left(\frac{\partial [F(T)]}{\partial s} \Big|_{s=0} \right)^2 \right\} dt \\ &= - \int_0^r \{ \mathbf{K}(\dot{\eta}, W) + g_{\dot{\eta}}(\dot{\eta}, D_W T)^2 \} dt \\ &= - \int_0^r \mathbf{K}(\dot{\eta}, W) dt. \end{aligned}$$

Since \mathbf{K} is continuous, letting $r \rightarrow 0$ yields $\mathbf{K}(v, w) \geq 0$.

The case of convex past capsules is shown in the same way, using the geodesic $\eta: [-r, 0] \rightarrow U_x$ with $\dot{\eta}(0) = v$. Alternatively, we can reduce the past case for L to the future case with respect to the reverse Lorentz–Finsler structure $\bar{L}(v) := L(-v)$, for the curvature bound $\mathbf{K} \geq 0$ (in timelike directions) for L is equivalent to that for \bar{L} (cf. [LMO1, Remark 8.14], [Oh2, §2.5]). \square

Remark 4.4 (Spacelike geodesics). Note that, by construction, $\dot{\sigma}_t(0) = W(t)$ is spacelike with respect to $g_{\dot{\eta}(t)}$. This is the reason why we introduced not only local timelike concavity but also local concavity in Definition 3.1, although it is desirable to deal only with causal curves. In fact, for spacelike geodesics, nothing can be said about their maximality or minimality; thus, it is not known how to consider spacelike geodesics in the synthetic setting like [BKR, EG, KS].

The proof of Theorem 1.1 is now complete: (I) \Rightarrow (II) by Theorem 3.6, (II) \Rightarrow (III) is obvious, (III) \Rightarrow (I) by Theorem 3.8, (II) \Rightarrow (IV) and (V) by Proposition 4.2, (IV) or (V) \Rightarrow (I) by Proposition 4.3.

5 Further problems

This section is devoted to discussions on some remaining open problems.

5.1 From concavity to the Berwald condition

In this subsection, we consider general (non-Berwald) Finsler spacetimes and explain difficulties in following the argument in [IL] in the positive definite case. Notice that Lemma 3.1 and Proposition 4.1 in [IL] can be generalized as follows.

Lemma 5.1 (cf. [IL, Lemma 3.1]). *Let (M, L) be a locally concave Finsler spacetime. Then, for every parallel vector field V along any nonconstant geodesic $\eta: [0, 1] \rightarrow M$ with $V(0) \in \Omega_{\eta(0)}$ (i.e., $D_{\dot{\eta}}^\eta V \equiv 0$), $L(V)$ is constant. In particular, $V(t) \in \Omega_{\eta(t)}$ for all $t \in [0, 1]$.*

Proof. Let J be the Jacobi field along η with $J(0) = 0$ and $D_{\dot{\eta}} J(0) = V(0)$. Observe that

$$D_{\dot{\eta}} D_{\dot{\eta}} J(0) = -R_{\dot{\eta}}(0) = 0.$$

Comparing this with $D_{\dot{\eta}}[tV(t)] = V(t)$ and $D_{\dot{\eta}} D_{\dot{\eta}}[tV(t)] = 0$, we find $J(t) = tV(t) + O(t^3)$. It follows that

$$F(J(t)) = tF(V(t)) + O(t^3) \quad \text{as } t \rightarrow 0.$$

Combining this with the concavity of $t \mapsto F(J(t))$ by Lemma 3.7, we obtain $F(J(t)) \leq tF(V(0))$. Hence, we have

$$F(V(t)) = \frac{1}{t}F(J(t)) + O(t^2) \leq F(V(0)) + O(t^2).$$

This implies

$$\frac{d}{dt}[F(V(t))]_{t=0} \leq 0,$$

and similarly $\frac{d}{dt}[F(V(t))] \leq 0$ for all $t \in (0, 1)$ such that $V(t)$ is timelike.

The same argument for the reverse Lorentz-Finsler structure $\bar{L}(v) := L(-v)$ shows that $\frac{d}{dt}[F(V(t))] \geq 0$ for all $t \in (0, 1]$ such that $V(t)$ is timelike. Indeed, $\bar{\eta}(t) := \eta(1 - t)$ is a geodesic for \bar{L} and $\frac{d}{dt}[\bar{F}(-V(t))] \geq 0$ follows with

$$\bar{F}(-V) := \sqrt{-2\bar{L}(-V)} = F(V).$$

Therefore, $F(V(t))$ is constant and $V(t)$ is timelike for all $t \in [0, 1]$. \square

Proposition 5.2 (cf. [11, Proposition 4.1]). *Let (M, L) be a Finsler spacetime, and suppose that there is a Lorentzian metric g on M which is preserved by every parallel transport along any geodesic. Then, g and L have the same geodesics and, in particular, L is of Berwald type.*

Proof. In this proof, we emphasize “ g -” for objects considered with respect to g ; otherwise, they are with respect to L .

Denote by ∇^g the g -covariant derivative. Fix $x \in M$ and, for $v \in T_x M$, we define

$$\kappa(v) := \nabla_t^g[\dot{\eta}_v](0) \in T_x M,$$

where η_v is the geodesic with $\dot{\eta}_v(0) = v$. We shall show that κ vanishes for all $v \in T_x M$, namely η_v is also a g -geodesic.

Given $w \in T_x M$, let W be the parallel vector field along η_v such that $W(0) = w$ (i.e., $D_{\dot{\eta}_v} W \equiv 0$). We claim that

$$\frac{d}{ds}[\kappa(v + sw)]_{s=0} = 2\nabla_t^g W(0). \quad (5.1)$$

The left-hand side can be understood in terms of local coordinates as

$$\sum_{i=1}^n \frac{d}{ds}[\kappa^i(v + sw)]_{s=0} \cdot \frac{\partial}{\partial x^i} \Big|_x \in T_x M.$$

To this end, we employ a normal coordinate system with respect to g around x (see, e.g., [10, Proposition 3.33]). Then, the g -Christoffel symbols at x vanish, thereby

$$\kappa(v) = \sum_{i=1}^n \ddot{\eta}_v^i(0) \frac{\partial}{\partial x^i} \Big|_x, \quad \nabla_t^g W(0) = \sum_{i=1}^n \dot{W}^i(0) \frac{\partial}{\partial x^i} \Big|_x.$$

This implies that

$$\frac{d}{ds}[\kappa(v + sw)]_{s=0} = \sum_{i=1}^n \frac{d}{ds}[\ddot{\eta}_{v+sw}^i(0)]_{s=0} \cdot \frac{\partial}{\partial x^i} \Big|_x = \sum_{i=1}^n \frac{d^2}{dt^2} \left[\frac{\partial}{\partial s} [\eta_{v+sw}^i(t)]_{s=0} \right]_{t=0} \cdot \frac{\partial}{\partial x^i} \Big|_x.$$

We set

$$J(t) := \sum_{i=1}^n \frac{\partial}{\partial s} [\eta_{v+sw}^i(t)]_{s=0} \cdot \frac{\partial}{\partial x^i} \Big|_x,$$

which is a Jacobi field along η_v such that $J(0) = 0$ and $D_{\dot{\eta}_v} J(0) = w$. It follows that $J(t) = tW(t) + O(t^3)$ (as in the proof of Lemma 5.1), and hence

$$\sum_{i=1}^n \ddot{J}^i(0) \frac{\partial}{\partial x^i} \Big|_x = \sum_{i=1}^n 2\dot{W}^i(0) \frac{\partial}{\partial x^i} \Big|_x = 2\nabla_t^g W(0).$$

This yields (5.1).

Now, for any parallel vector fields V, W along η_v , we infer from the hypothesis that

$$0 = \frac{d}{dt} [g(V, W)] = g(\nabla_t^g V, W) + g(V, \nabla_t^g W). \quad (5.2)$$

On the one hand, plugging $V = W = \dot{\eta}_{v+sw}$ into (5.2) at $t = 0$ implies $g(\kappa(v + sw), v + sw) = 0$. Combining this with (5.1), we find, for W as in (5.1),

$$0 = \frac{d}{ds} [g(\kappa(v + sw), v + sw)]_{s=0} = 2g(\nabla_t^g W(0), v) + g(\kappa(v), w).$$

On the other hand, (5.2) with $V = \dot{\eta}_v$ and W as in (5.1) at $t = 0$ shows

$$g(\kappa(v), w) + g(v, \nabla_t^g W(0)) = 0.$$

Comparing these equations, we obtain $g(\kappa(v), w) = 0$. Since $w \in T_x M$ was arbitrary, $\kappa(v) = 0$ holds. Therefore, g and L have the same geodesics. This implies that the Chern connection of L coincides with the Levi-Civita connection of g , thereby L is of Berwald type. \square

Remark 5.3 (Canonical Lorentzian metrics). In order to build a bridge between Lemma 5.1 and Proposition 5.2, in the situation of Lemma 5.1, we need to construct a Lorentzian metric g satisfying the hypothesis in Proposition 5.2. In the positive definite case, it was done in [IL, Proposition 4.2] by using a Riemannian metric “canonically” associated with a Finsler metric. Such a construction is not known in Lorentz–Finsler geometry; this is indeed a major obstacle in the metrization problem (recall Remark 2.4). It is sufficient to find a “canonical” way of choosing a finite measure on each tangent space $T_x M$, as taking an average of g_v in $v \in T_x M$ with that measure provides a Lorentzian metric (in the positive definite case, the uniform measure on the unit ball or sphere plays a role). However, the noncompactness of the isometry group of Minkowski metrics arises as an essential difficulty in such a construction.

5.2 Concavity and the variance functional

In the positive definite case, there is an interesting nonsmooth characterization of Busemann’s convexity in terms of optimal transport theory. We refer to [Vi] for the basic knowledge of optimal transport theory.

Let (X, d) be a complete geodesic space. Given a Borel probability measure $\mu \in \mathcal{P}_2(X)$ of finite second moment, its *variance* is defined by

$$\text{var}(\mu) := \inf_{x \in X} \int_X d^2(x, y) \mu(dy).$$

A point $x \in X$ achieving the above infimum is called a *barycenter* (also called *center of mass* or *Fréchet mean*) of μ . If (X, d) is globally convex, then the set of barycenters of μ is a convex set. In the case where (X, d) is a complete CAT(0)-space, any μ admits a unique barycenter.

Proposition 5.4 (Kim–Pass [KP]). *A complete separable geodesic space (X, d) is globally convex if and only if $\sqrt{\text{var}}$ is convex along geodesics in the L^2 -Wasserstein space $(\mathcal{P}_2(X), W_2)$.*

Proof. To be precise, it was shown in [KP, Proposition 2.1] that d^2 is convex if and only if var is convex. For completeness, we give a proof along the same lines as [KP].

In the “if” part, given minimizing geodesics $\gamma_1, \gamma_2: [0, 1] \rightarrow X$, with $\gamma_1(0) = \gamma_2(0)$, consider probability measures $\mu_t := (\delta_{\gamma_1(t)} + \delta_{\gamma_2(t)})/2$. Then, $(\mu_t)_{t \in [0, 1]}$ is a W_2 -geodesic and

$$\text{var}(\mu_t) = \frac{1}{4} d^2(\gamma_1(t), \gamma_2(t))$$

for all $t \in [0, 1]$ (any midpoint of $\gamma_1(t)$ and $\gamma_2(t)$ is a barycenter of μ_t). Hence, the convexity of $\sqrt{\text{var}(\mu_t)}$ implies that $t \mapsto d(\gamma_1(t), \gamma_2(t))$ is convex.

For general minimizing geodesics $\gamma_1, \gamma_2: [0, 1] \rightarrow X$, let $\eta: [0, 1] \rightarrow X$ be a minimizing geodesic from $\gamma_1(0)$ to $\gamma_2(1)$. Then, we deduce from the triangle inequality and the above estimate that

$$\begin{aligned} d(\gamma_1(t), \gamma_2(t)) &\leq d(\gamma_1(t), \eta(t)) + d(\eta(t), \gamma_2(t)) \\ &\leq t d(\gamma_1(1), \gamma_2(1)) + (1 - t) d(\gamma_1(0), \gamma_2(0)). \end{aligned}$$

Thus, (X, d) is globally convex.

To see the “only if” part, let $\mu_t = (e_t)_* \Pi$ be a W_2 -geodesic induced from a probability measure Π on the set $\Gamma(X)$ of minimizing geodesics $\eta: [0, 1] \rightarrow X$, where $e_t(\eta) := \eta(t)$ and $(e_t)_* \Pi$ denotes the push-forward of Π by e_t (see [Li, Theorem 6]). For arbitrary $\varepsilon > 0$, take $x_i \in X$ such that

$$\left(\int_X d^2(x_i, y) \mu_i(dy) \right)^{1/2} \leq \sqrt{\text{var}(\mu_i)} + \varepsilon, \quad i = 0, 1.$$

Let $\gamma: [0, 1] \rightarrow X$ be the (unique) minimizing geodesic from x_0 to x_1 . Then, it follows from the global convexity of (X, d) that

$$\begin{aligned} \sqrt{\text{var}(\mu_t)} &\leq \left(\int_X d^2(\gamma(t), y) \mu_t(dy) \right)^{1/2} = \left(\int_{\Gamma(X)} d^2(\gamma(t), \eta(t)) \Pi(d\eta) \right)^{1/2} \\ &\leq \left(\int_{\Gamma(X)} \left((1 - t) d(\gamma(0), \eta(0)) + t d(\gamma(1), \eta(1)) \right)^2 \Pi(d\eta) \right)^{1/2} \\ &\leq (1 - t) \left(\int_{\Gamma(X)} d^2(\gamma(0), \eta(0)) \Pi(d\eta) \right)^{1/2} + t \left(\int_{\Gamma(X)} d^2(\gamma(1), \eta(1)) \Pi(d\eta) \right)^{1/2} \\ &= (1 - t) \left(\int_X d^2(x_0, y) \mu_0(dy) \right)^{1/2} + t \left(\int_X d^2(x_1, y) \mu_1(dy) \right)^{1/2} \\ &\leq (1 - t) \sqrt{\text{var}(\mu_0)} + t \sqrt{\text{var}(\mu_1)} + \varepsilon. \end{aligned}$$

Letting $\varepsilon \rightarrow 0$ shows that $t \mapsto \sqrt{\text{var}(\mu_t)}$ is convex. \square

Remark 5.5 (Variance and barycenter). To formulate a Lorentzian counterpart to Proposition 5.4, we need an appropriate notion of variance, which is an open question. This is related to another open question: how to define the barycenter of a probability measure in a Lorentzian or Finsler spacetime? Even in the Minkowski space (\mathbf{R}^n, L) , no reasonable characterization of the linear average $\int_{\mathbf{R}^n} x \mu(dx)$ in a synthetic/metric way seems to be known. These notions will be helpful to develop probability theory and statistics on spacetimes.

Acknowledgements This research was supported in part by the Austrian Science Fund (FWF) [Grants DOI [10.55776/PAT1996423](#) and [10.55776/EF6](#)]. SO was supported by the JSPS Grant-in-Aid for Scientific Research (KAKENHI) 22H04942, 24K00523. FR acknowledges the support of the European Union - NextGenerationEU, in the framework of the PRIN Project ‘Contemporary perspectives on geometry and gravity’ (code 2022JJ8KER – CUP G53D23001810006). The views and opinions expressed are solely those of the authors and do not necessarily reflect those of the European Union, nor can the European Union be held responsible for them.

For open access purposes, the authors have applied a [Creative Commons Attribution 4.0 International](#) license to any author-accepted manuscript version arising from this submission.

References

- [AB] S. B. Alexander and R. L. Bishop, Lorentz and semi-Riemannian spaces with Alexandrov curvature bounds. *Comm. Anal. Geom.* **16** (2008), 251–282.
- [AKP] S. Alexander, V. Kapovitch and A. Petrunin, *Alexandrov geometry—foundations*. American Mathematical Society, Providence, RI, 2024.
- [BCS] D. Bao, S.-S. Chern and Z. Shen, *An introduction to Riemann–Finsler geometry*. Springer-Verlag, New York, 2000.
- [Be] J. K. Beem, Indefinite Finsler spaces and timelike spaces. *Can. J. Math.* **22** (1970), 1035–1039.
- [BEE] J. K. Beem, P. E. Ehrlich and K. L. Easley, *Global Lorentzian geometry*. Marcel Dekker Inc., New York, 1996.
- [BKOR] T. Beran, M. Kunzinger, A. Ohanyan and F. Rott, The equivalence of smooth and synthetic notions of timelike sectional curvature bounds. Preprint (2024), [arXiv:2403.18077](#)
- [BKR] T. Beran, M. Kunzinger and F. Rott, On curvature bounds in Lorentzian length spaces. *J. Lond. Math. Soc. (2)* **110** (2024), Paper No. e12971.
- [BORS] T. Beran, A. Ohanyan, F. Rott and D. A. Solis, The splitting theorem for globally hyperbolic Lorentzian length spaces with non-negative timelike curvature. *Lett. Math. Phys.* **113** (2023), Paper No. 48.
- [BS] T. Beran and C. Sämann, Hyperbolic angles in Lorentzian length spaces and timelike curvature bounds. *J. Lond. Math. Soc. (2)* **107** (2023), 1823–1880.

- [BO] M. Braun and S. Ohta, Optimal transport and timelike lower Ricci curvature bounds on Finsler spacetimes. *Trans. Amer. Math. Soc.* **377** (2024), 3529–3576.
- [BH] M. R. Bridson and A. Haefliger, Metric spaces of non-positive curvature. *Grundlehren der mathematischen Wissenschaften*, **319**. Springer-Verlag, Berlin, 1999.
- [BBI] D. Burago, Y. Burago and S. Ivanov, A course in metric geometry. American Mathematical Society, Providence, RI, 2001.
- [Bu] H. Busemann, Spaces with non-positive curvature. *Acta Math.* **80** (1948), 259–310.
- [COO] E. Caponio, A. Ohanyan and S. Ohta, Splitting theorems for weighted Finsler spacetimes via the p -d’Alembertian: beyond the Berwald case. Preprint (2024), [arXiv:2412.20783](#)
- [CM] F. Cavalletti and A. Mondino, Optimal transport in Lorentzian synthetic spaces, synthetic timelike Ricci curvature lower bounds and applications. *Camb. J. Math.* **12** (2024), 417–534.
- [EG] D. Erös and S. Gieger, A synthetic Lorentzian Cartan–Hadamard theorem. Preprint (2025), [arXiv:2506.22197](#)
- [FG] T. Fujioka and S. Gu, Topological regularity of Busemann spaces of nonpositive curvature. Preprint (2025), [arXiv:2504.14455](#)
- [FHPV] A. Fuster, S. Heefer, C. Pfeifer and N. Voicu, On the non metrizable of Berwald Finsler spacetimes. *Universe* **6** (2020), no. 5, Paper No. 64.
- [HY] B. Han and L. Yin, On the structure of Busemann spaces with non-negative curvature. Preprint (2025), [arXiv:2508.12348](#)
- [Ha] S. G. Harris, A triangle comparison theorem for Lorentz manifolds. *Indiana Univ. Math. J.* **31** (1982), no. 3, 289–308.
- [IL] S. Ivanov and A. Lytchak, Rigidity of Busemann convex Finsler metrics. *Comment. Math. Helv.* **94** (2019), 855–868.
- [KP] Y.-H. Kim and B. Pass, Nonpositive curvature, the variance functional, and the Wasserstein barycenter. *Proc. Amer. Math. Soc.* **148** (2020), 1745–1756.
- [KK] A. Kristály and L. Kozma, Metric characterization of Berwald spaces of non-positive flag curvature. *J. Geom. Phys.* **56** (2006), 1257–1270.
- [KVK] A. Kristály, C. Varga and L. Kozma, The dispersing of geodesics in Berwald spaces of non-positive flag curvature. *Houston J. Math.* **30** (2004), 413–420.
- [KS] M. Kunzinger and C. Sämann, Lorentzian length spaces. *Ann. Global Anal. Geom.* **54** (2018), 399–447.
- [Li] S. Lisini, Characterization of absolutely continuous curves in Wasserstein spaces. *Calc. Var. Partial Differential Equations* **28** (2007), 85–120.

- [LV] J. Lott and C. Villani, Ricci curvature for metric-measure spaces via optimal transport. *Ann. of Math. (2)* **169** (2009), 903–991.
- [LMO1] Y. Lu, E. Minguzzi and S. Ohta, Geometry of weighted Lorentz–Finsler manifolds I: Singularity theorems. *J. Lond. Math. Soc.* **104** (2021), 362–393.
- [LMO2] Y. Lu, E. Minguzzi and S. Ohta, Comparison theorems on weighted Finsler manifolds and spacetimes with ε -range. *Anal. Geom. Metr. Spaces* **10** (2022), 1–30.
- [LMO3] Y. Lu, E. Minguzzi and S. Ohta, Geometry of weighted Lorentz–Finsler manifolds II: A splitting theorem. *Internat. J. Math.* **34** (2023), Paper No. 2350002.
- [Mc] R. J. McCann, Displacement convexity of Boltzmann’s entropy characterizes the strong energy condition from general relativity. *Camb. J. Math.* **8** (2020), 609–681.
- [Mi1] E. Minguzzi, Convex neighborhoods for Lipschitz connections and sprays. *Monatsh. Math.* **177** (2015), 569–625.
- [Mi2] E. Minguzzi, Raychaudhuri equation and singularity theorems in Finsler spacetimes. *Class. Quantum Grav.* **32**, 185008 (2015).
- [MS] A. Mondino and S. Suhr, An optimal transport formulation of the Einstein equations of general relativity. *J. Eur. Math. Soc. (JEMS)* **25** (2023), 933–994.
- [Oh1] S. Ohta, On the measure contraction property of metric measure spaces. *Comment. Math. Helv.* **82** (2007), 805–828.
- [Oh2] S. Ohta, Comparison Finsler geometry. *Springer Monographs in Mathematics*. Springer, Cham, 2021.
- [ON] B. O’Neill, *Semi-Riemannian geometry: With applications to relativity*. Academic Press, Inc., New York, 1983.
- [St1] K.-T. Sturm, On the geometry of metric measure spaces. I. *Acta Math.* **196** (2006), 65–131.
- [St2] K.-T. Sturm, On the geometry of metric measure spaces. II. *Acta Math.* **196** (2006), 133–177.
- [Sz] Z. I. Szabó, Positive definite Berwald spaces. Structure theorems on Berwald spaces. *Tensor (N.S.)* **35** (1981), 25–39.
- [Vi] C. Villani, *Optimal transport, old and new*. Springer-Verlag, Berlin, 2009.